


LPP polyclonal antibody

Catalog # PAB17854 Size 100 ug

Applications



Western Blot (Cell lysate)

LPP polyclonal antibody (Cat # PAB17854) (0.1 ug/mL) staining of HepG2 lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Specification

Product Description	Goat polyclonal antibody raised against synthetic peptide of LPP.
Immunogen	A synthetic peptide corresponding to amino acids at internal region of human LPP.
Sequence	C-RNDSPTYGQQGHP
Host	Goat
Theoretical MW (kDa)	65.7
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Concentration	0.5 mg/mL
Recommend Usage	ELISA (1:16000) Western Blot (0.1-0.3 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 0.5 mg/mL Tris saline, pH 7.3 (0.02% sodium azide, 0.5% BSA)

Storage Instruction

Store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

LPP polyclonal antibody (Cat # PAB17854) (0.1 ug/mL) staining of HepG2 lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — LPP

Entrez GeneID [4026](#)

Protein Accession# [NP_005569.1](#)

Gene Name LPP

Gene Alias -

Gene Description LIM domain containing preferred translocation partner in lipoma

Omim ID [600700 601626](#)

Gene Ontology [Hyperlink](#)

Other Designations LIM domain-containing preferred translocation partner in lipoma|LIM protein|Lipoma-preferred partner

Disease

- [Autoimmune Diseases](#)

- [Celiac Disease](#)

- [Diabetes Mellitus](#)

- [Genetic Predisposition to Disease](#)

- [Melanoma](#)
- [Tobacco Use Disorder](#)
- [Vitiligo](#)