


PRKDC polyclonal antibody

Catalog # PAB18344 Size 100 ug

Applications



Immunofluorescence

Immunofluorescence analysis of HUVEC cells, treated with serum (20 %, 30 mins), using PRKDC polyclonal antibody (Cat # PAB18344).
Peptide "+" means "peptide blocking".

Specification

Product Description Rabbit polyclonal antibody raised against synthetic peptide of PRKDC.

Immunogen A synthetic peptide corresponding to human PRKDC.

Host Rabbit

Reactivity Human

Specificity This antibody is specific to PRKDC.

Form Liquid

Purification Affinity purification

Concentration 1 mg/mL

Recommend Usage
Immunofluorescence (1:500-1:1000)
ELISA (1:20000)
The optimal working dilution should be determined by the end user.

Storage Buffer In PBS, 150mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)

Storage Instruction

Store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunofluorescence

Immunofluorescence analysis of HUVEC cells, treated with serum (20 %, 30 mins), using PRKDC polyclonal antibody (Cat # PAB18344).
Peptide "+" means "peptide blocking".

- Enzyme-linked Immunoabsorbent Assay

Gene Info — PRKDC

Entrez GeneID[5591](#)**Protein Accession#**[P78527](#)**Gene Name**

PRKDC

Gene Alias

DNA-PKcs, DNAPK, DNP1, HYRC, HYRC1, XRCC7, p350

Gene Description

protein kinase, DNA-activated, catalytic polypeptide

Omim ID[600899](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

The PRKDC gene encodes the catalytic subunit of a nuclear DNA-dependent serine/threonine protein kinase (DNA-PK). The second component is the autoimmune antigen Ku (MIM 152690), which is encoded by the G22P1 gene on chromosome 22q. On its own, the catalytic subunit of DNA-PK is inactive and relies on the G22P1 component to direct it to the DNA and trigger its kinase activity; PRKDC must be bound to DNA to express its catalytic properties.[supplied by OMIM]

Other Designations

hyper-radiosensitivity of murine scid mutation, complementing 1

Publication Reference

- [Frameshift mutation in PRKDC, the gene for DNA-PKcs, in the DNA repair-defective, human, glioma-derived cell line M059J.](#)

Anderson CW, Dunn JJ, Freimuth PI, Galloway AM, Allalunis-Turner MJ.

Radiation Research 2001 Jul; 156(1):2.

Pathway

- [Cell cycle](#)
- [Non-homologous end-joining](#)

Disease

- [Adenocarcinoma](#)
- [Ataxia telangiectasia](#)
- [Bone Neoplasms](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Central Nervous System Neoplasms](#)
- [DNA Damage](#)
- [Genetic Predisposition to Disease](#)
- [Glioma](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Neoplasms](#)
- [Lung Neoplasms](#)

- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Sclerosis](#)
- [Neoplasms](#)
- [Occupational Diseases](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Thyroid Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)