


RPS6KB1 polyclonal antibody

Catalog # PAB18495 Size 100 ug


Applications



Western Blot (Cell lysate)

Western blot analysis of extracts from HeLa cells, untreated or treated with insulin (10 U/mL, 5 min), using RPS6KB1 polyclonal antibody (Cat # PAB18495).

Peptide "+" means "peptide blocking".



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using RPS6KB1 polyclonal antibody (Cat # PAB18495).

Peptide "+" means "peptide blocking".

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of RPS6KB1.
Immunogen	A synthetic peptide corresponding to residues surrounding S411 of human RPS6KB1.
Host	Rabbit
Reactivity	Human, Mouse, Rat
Specificity	This antibody is specific to RPS6KB1.
Form	Liquid

Purification	Affinity purification
Concentration	1 mg/mL
Recommend Usage	Western Blot (1:500-1:1000) Immunohistochemistry (1:50-1:100) ELISA (1:10000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of extracts from HeLa cells, untreated or treated with insulin (10 U/mL, 5 min), using RPS6KB1 polyclonal antibody (Cat # PAB18495).

Peptide "+" means "peptide blocking".

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using RPS6KB1 polyclonal antibody (Cat # PAB18495).

Peptide "+" means "peptide blocking".

- Enzyme-linked Immunoabsorbent Assay

Gene Info — RPS6KB1

Entrez GeneID	6198
Protein Accession#	P23443
Gene Name	RPS6KB1
Gene Alias	PS6K, S6K, S6K1, STK14A, p70(S6K)-alpha, p70-S6K, p70-alpha
Gene Description	ribosomal protein S6 kinase, 70kDa, polypeptide 1
Omim ID	608938

Gene Ontology

[Hyperlink](#)

Gene Summary

This gene encodes a member of the RSK (ribosomal S6 kinase) family of serine/threonine kinase s. This kinase contains 2 non-identical kinase catalytic domains and phosphorylates several residues of the S6 ribosomal protein. The kinase activity of this protein leads to an increase in protein synthesis and cell proliferation. Amplification of the region of DNA encoding this gene and overexpression of this kinase are seen in some breast cancer cell lines. Alternate translational start sites have been described and alternate transcriptional splice variants have been observed but have not been thoroughly characterized. [provided by RefSeq]

Other Designations

p70 S6 kinase, alpha 1|p70 S6 kinase, alpha 2|ribosomal protein S6 kinase, 70kD, polypeptide 1|serine/threonine kinase 14 alpha

Publication Reference

- [Glycogen synthase kinase-3 interacts with and phosphorylates estrogen receptor alpha and is involved in the regulation of receptor activity.](#)

Medunjanin S, Hermani A, De Servi B, Grisouard J, Rincke G, Mayer D.

The Journal of Biological Chemistry 2005 Aug; 280(38):33006.

- [Ligand-independent interactions of p160/steroid receptor coactivators and CREB-binding protein \(CBP\) with estrogen receptor-alpha: regulation by phosphorylation sites in the A/B region depends on other receptor domains.](#)

Dutertre M, Smith CL.

Molecular Endocrinology (Baltimore, Md.) 2003 Apr; 17(7):1296.

- [Control of p70 ribosomal protein S6 kinase and acetyl-CoA carboxylase by AMP-activated protein kinase and protein phosphatases in isolated hepatocytes.](#)

Krause U, Bertrand L, Hue L.

European Journal of Biochemistry 2002 Aug; 269(15):3751.

- [A new role for the p85-phosphatidylinositol 3-kinase regulatory subunit linking FRAP to p70 S6 kinase activation.](#)

Gonzalez-Garcia A, Garrido E, Hernandez C, Alvarez B, Jimenez C, Cantrell DA, Pullen N, Carrera AC.

Journal of Biological Chemistry 2002 Jan; 277(2):1500.

- [Activation of estrogen receptor alpha by S118 phosphorylation involves a ligand-dependent interaction with TFIIF and participation of CDK7.](#)

Chen D, Riedl T, Washbrook E, Pace PE, Coombes RC, Egly JM, Ali S.

Molecular Cell 2000 Jul; 6(1):127.

Pathway

- [Acute myeloid leukemia](#)
- [ErbB signaling pathway](#)
- [Fc gamma R-mediated phagocytosis](#)
- [Insulin signaling pathway](#)
- [mTOR signaling pathway](#)
- [TGF-beta signaling pathway](#)

Disease

- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Head and Neck Neoplasms](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)