


SP1 polyclonal antibody

Catalog # PAB18500 Size 100 ug


Applications



Western Blot (Cell lysate)

Western blot analysis of extracts from HT-29 cells, using SP1 polyclonal antibody (Cat # PAB18500).

Peptide "+" means "peptide blocking".



Immunofluorescence

Immunofluorescence analysis of HepG2 cells, using SP1 polyclonal antibody (Cat # PAB18500).

Peptide "+" means "peptide blocking".

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of SP1.
Immunogen	A synthetic peptide corresponding to residues surrounding T739 of human SP1.
Host	Rabbit
Reactivity	Human, Mouse, Rat
Specificity	This antibody is specific to SP1.
Form	Liquid

Purification	Affinity purification
Concentration	1 mg/mL
Recommend Usage	Western Blot (1:500-1:1000) Immunohistochemistry (1:50-1:100) Immunofluorescence (1:500-1:1000) ELISA (1:20000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of extracts from HT-29 cells, using SP1 polyclonal antibody (Cat # PAB18500).
Peptide "+" means "peptide blocking".

- Immunohistochemistry

- Immunofluorescence

Immunofluorescence analysis of HepG2 cells, using SP1 polyclonal antibody (Cat # PAB18500).
Peptide "+" means "peptide blocking".

- Enzyme-linked Immunoabsorbent Assay

Gene Info — SP1

Entrez GeneID	6667
Protein Accession#	P08047
Gene Name	SP1
Gene Alias	-
Gene Description	Sp1 transcription factor

Omim ID	189906
Gene Ontology	Hyperlink
Other Designations	specificity protein 1

Publication Reference

- [The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection \(MGC\).](#)

Gerhard DS, Wagner L, Feingold EA, Shenmen CM, Grouse LH, Schuler G, Klein SL, Old S, Rasooly R, Good P, Guyer M, Peck AM, Derge JG, Lipman D, Collins FS, Jang W, Sherry S, Feolo M, Misquitta L, Lee E, Rotmistrovsky K, Greenhut SF, Schaefer CF, Buetow K, Bonner TI, Haussler D, Kent J, Kiekhaus M, Furey T, Brent M, Prange C, Schreiber K, Shapiro N, Bhat NK, Hopkins RF, Hsie F, Driscoll T, Soares MB, Casavant TL, Scheetz TE, Brown-stein MJ, Usdin TB, Toshiyuki S, Carninci P, Piao Y, Dudekula DB, K

Genome Research 2004 Oct; 14(10B):2121.

- [Multiple elements regulate nuclear/cytoplasmic shuttling of FOXO1: characterization of phosphorylation- and 14-3-3-dependent and -independent mechanisms.](#)

Zhao X, Gan L, Pan H, Kan D, Majeski M, Adam SA, Unterman TG.

The Biochemical Journal 2004 Mar; 378(Pt 3):839.

- [Two novel phosphorylation sites on FKHR that are critical for its nuclear exclusion.](#)

Rena G, Woods YL, Prescott AR, Peggie M, Unterman TG, Williams MR, Cohen P.

The EMBO Journal 2002 May; 21(9):2263.

- [Roles of the forkhead in rhabdomyosarcoma \(FKHR\) phosphorylation sites in regulating 14-3-3 binding, transactivation and nuclear targetting.](#)

Rena G, Prescott AR, Guo S, Cohen P, Unterman TG.

The Biochemical Journal 2001 Mar; 354(Pt 3):605.

- [Heterogeneous Sp1 mRNAs in human HepG2 cells include a product of homotypic trans-splicing.](#)

Takahara T, Kanazu SI, Yanagisawa S, Akanuma H.

The Journal of Biological Chemistry 2000 Dec; 275(48):38067.

Pathway

- [TGF-beta signaling pathway](#)

Disease

- [Head and Neck Neoplasms](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)