


POU3F2 polyclonal antibody

Catalog # PAB18579

Size 100 ug

Applications



Western Blot (Tissue lysate)

POU3F2 polyclonal antibody (Cat # PAB18579) (0.1 ug/mL) staining of mouse brain lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Specification

Product Description	Goat polyclonal antibody raised against synthetic peptide of POU3F2.
Immunogen	A synthetic peptide corresponding to amino acids at N-terminus of human POU3F2.
Sequence	C-AQSLVQGDY GALQSN
Host	Goat
Theoretical MW (kDa)	45
Reactivity	Human, Mouse, Rat
Form	Liquid
Purification	Antigen affinity purification
Concentration	0.5 mg/mL
Recommend Usage	ELISA (1:8000) Western Blot (0.1-0.3 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In 0.5 mg/mL Tris saline, pH 7.3 (0.02% sodium azide, 0.5% BSA)

Storage Instruction

Store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Tissue lysate)

POU3F2 polyclonal antibody (Cat # PAB18579) (0.1 ug/mL) staining of mouse brain lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — POU3F2

Entrez GeneID[5454](#)**Protein Accession#**[NP_005595.2](#)**Gene Name**

POU3F2

Gene Alias

BRN2, OCT7, OTF7, POUF3

Gene Description

POU class 3 homeobox 2

Omim ID[600494](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

POU3F2 belongs to a large family of transcription factors that bind to the octameric DNA sequence ATGCAAAT. Most of these proteins share a highly homologous region, referred to as the POU domain, that occurs in several mammalian transcription factors, including the octamer-binding proteins Oct1 (POU2F1; MIM 164175) and Oct2 (POU2F2; MIM 164176) and the pituitary protein Pit 1 (PIT1; MIM 173110). Class III POU genes are expressed predominantly in the central nervous system (CNS). It is likely that CNS-specific transcription factors such as these play an important role in mammalian neurogenesis by regulating their diverse patterns of gene expression (Schreiber et al., 1993 [PubMed 8441633]; Atanasoski et al., 1995 [PubMed 7601453]).[supplied by OMIM]

Other Designations

POU domain, class 3, transcription factor 2

Disease

- [Huntington disease](#)
- [Schizophrenia](#)