

FBXO40 polyclonal antibody

Catalog # PAB18728

Size 100 ug

Specification

| | |
|----------------------------|--|
| Product Description | Goat polyclonal antibody raised against synthetic peptide of FBXO40. |
| Immunogen | A synthetic peptide corresponding to amino acids at internal region of human FBXO40. |
| Sequence | C-KEPQENQKQQDVRT |
| Host | Goat |
| Form | Liquid |
| Purification | Antigen affinity purification |
| Concentration | 0.5 mg/mL |
| Recommend Usage | ELISA (1:128000) The optimal working dilution should be determined by the end user. |
| Storage Buffer | In Tris saline, pH 7.3 (0.02% sodium azide, 0.5% BSA) |
| Storage Instruction | Store at -20°C. Aliquot to avoid repeated freezing and thawing. |
| Note | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |

Applications

- Enzyme-linked Immunoabsorbent Assay

Gene Info — FBXO40

Entrez GeneID [51725](#)

| | |
|--------------------|--|
| Protein Accession# | NP_057382.2 |
| Gene Name | FBXO40 |
| Gene Alias | FBX40, KIAA1195, MGC129902, MGC129903 |
| Gene Description | F-box protein 40 |
| Omim ID | 609107 |
| Gene Ontology | Hyperlink |
| Gene Summary | Members of the F-box protein family, such as FBXO40, are characterized by an approximately 40 -amino acid F-box motif. SCF complexes, formed by SKP1 (MIM 601434), cullin (see CUL1; MIM 603134), and F-box proteins, act as protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact with ubiquitination targets through other protein interaction domains (Jin et al., 2004 [PubMed 15520277]).[supplied by OMIM] |
| Other Designations | muscle disease-related protein |