


GHITM polyclonal antibody

Catalog # PAB18758 Size 100 ug

Applications



Western Blot (Cell lysate)

GHITM polyclonal antibody (Cat # PAB18758) (0.3 ug/mL) staining of HeLa cell lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Specification

Product Description	Goat polyclonal antibody raised against synthetic peptide of GHITM.
Immunogen	A synthetic peptide corresponding to amino acids at internal region of human GHITM.
Sequence	C-PSREYATKTRIGIR
Host	Goat
Theoretical MW (kDa)	35
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Concentration	0.5 mg/mL
Recommend Usage	ELISA (1:8000) Western Blot (0.1-0.5 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In Tris saline, pH 7.3 (0.02% sodium azide, 0.5% BSA)

Storage Instruction

Store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

GHITM polyclonal antibody (Cat # PAB18758) (0.3 ug/mL) staining of HeLa cell lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — GHITM

Entrez GeneID[27069](#)**Protein Accession#**[NP_055209.2](#)**Gene Name**

GHITM

Gene Alias

DERP2, DKFZp566C0746, FLJ26584, HSPC282, MICS1, My021, PTD010, TMBIM5

Gene Description

growth hormone inducible transmembrane protein

Gene Ontology[Hyperlink](#)**Other Designations**

OTTHUMP00000019998|dermal papilla-derived protein 2|mitochondrial morphology and cristae structure 1|transmembrane BAX inhibitor motif containing 5

Disease

- [Alzheimer Disease](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)