


RELA polyclonal antibody

Catalog # PAB18764 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of formalin-fixed paraffin-embedded human tonsil tissue showing nuclear staining with RELA polyclonal antibody (Cat # PAB18764) at 1 : 100 dilution.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of RELA.
Immunogen	A synthetic peptide corresponding to 14 amino acids at N-terminus of human RELA.
Host	Rabbit
Reactivity	Human
Form	Liquid
Recommend Usage	ELISA (1:20000-1:80000) Western Blot (1:200-1:1000) Immunohistochemistry (1:50-1:250) The optimal working dilution should be determined by the end user.
Storage Buffer	In buffer containing 0.02% sodium azide
Storage Instruction	Store at 4°C for three months. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of formalin-fixed paraffin-embedded human tonsil tissue showing nuclear staining with RELA polyclonal antibody (Cat # PAB18764) at 1 : 100 dilution.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — RELA

Entrez GeneID [5970](#)

Protein Accession# [NM_021975](#)

Gene Name RELA

Gene Alias MGC131774, NFKB3, p65

Gene Description v-rel reticuloendotheliosis viral oncogene homolog A (avian)

Omim ID [164014](#)

Gene Ontology [Hyperlink](#)

Gene Summary NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL (MIM 164910), RELA, or RELB (MIM 604758) to form the NFKB complex. The p50 (NFKB1)/p65 (RELA) heterodimer is the most abundant form of NFKB. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, MIM 164008 or NFKBIB, MIM 604495), which inactivate NFKB by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, MIM 600664, or IKBKB, MIM 603258) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NFKB complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).[supplied by OMIM]

Other Designations nuclear factor of kappa light polypeptide gene enhancer in B-cells 3|v-rel avian reticuloendotheliosis viral oncogene homolog A (nuclear factor of kappa light polypeptide gene enhancer in B-cells 3 (p65))|v-rel reticuloendotheliosis viral oncogene homolog

Pathway

- [Acute myeloid leukemia](#)

- [Adipocytokine signaling pathway](#)
- [Apoptosis](#)
- [B cell receptor signaling pathway](#)
- [Chemokine signaling pathway](#)
- [Chronic myeloid leukemia](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Small cell lung cancer](#)
- [T cell receptor signaling pathway](#)
- [Toll-like receptor signaling pathway](#)

Disease

- [Arthritis](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hematologic Diseases](#)

- [HIV Infections](#)
- [Hodgkin Disease](#)
- [Liver Cirrhosis](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Myeloma](#)
- [Occupational Diseases](#)
- [Testicular Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)