


GAD2 polyclonal antibody

Catalog # PAB19504 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded rat brain tissue section with GAD2 polyclonal antibody (Cat # PAB19504).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of GAD2.
Immunogen	A synthetic peptide corresponding to amino acids at N-terminus of human GAD2.
Host	Rabbit
Reactivity	Rat
Specificity	It identical to the related mouse and rat sequence.
Form	Lyophilized
Purification	Immunoaffinity purification
Isotype	IgG
Recommend Usage	Western Blot (1 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ (5 mg BSA, 0.05 mg sodium azide, 0.05 mg Thimerosal)

Storage Instruction

Store at -20°C on dry atmosphere.
After reconstitution with 200 uL of deionized water and concentration will be 500 ug/mL, store at -20°C or lower.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide and thimerosal: POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded rat brain tissue section with GAD2 polyclonal antibody (Cat # PAB19504).

Gene Info — GAD2

Entrez GeneID

[2572](#)

Gene Name

GAD2

Gene Alias

GAD65, MGC161605, MGC161607

Gene Description

glutamate decarboxylase 2 (pancreatic islets and brain, 65kDa)

Omim ID

[138275](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

This gene encodes one of several forms of glutamic acid decarboxylase, identified as a major autoantigen in insulin-dependent diabetes. The enzyme encoded is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. A pathogenic role for this enzyme has been identified in the human pancreas since it has been identified as an autoantibody and an autoreactive T cell target in insulin-dependent diabetes. This gene may also play a role in the stiff man syndrome. Alternative splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq]

Other Designations

Glutamate decarboxylase-2 (pancreas)|OTTHUMP00000019335|OTTHUMP00000019336|glutamate decarboxylase 2|glutamate decarboxylase 2 (pancreatic islets and brain, 65kD)

Pathway

- [Alanine](#)

- [beta-Alanine metabolism](#)
- [Butanoate metabolism](#)
- [Metabolic pathways](#)
- [Taurine and hypotaurine metabolism](#)
- [Type I diabetes mellitus](#)

Disease

- [Alcoholism](#)
- [Alzheimer disease](#)
- [Anorexia Nervosa](#)
- [Anxiety Disorders](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Birth Weight](#)
- [Body Weight](#)
- [Bulimia](#)
- [Cardiovascular Diseases](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Cognition](#)
- [Depressive Disorder](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Disease Models](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)

- [Hyperandrogenism](#)
- [Hypercholesterolemia](#)
- [Hyperglycemia](#)
- [Insulin Resistance](#)
- [Mental Disorders](#)
- [Metabolic Syndrome X](#)
- [NARP](#)
- [Neoplasms](#)
- [Neurotic Disorders](#)
- [Obesity](#)
- [Osteoporosis](#)
- [Psychiatric Status Rating Scales](#)
- [Puberty](#)
- [Schizophrenia](#)
- [Schizophrenic Psychology](#)
- [Tobacco Use Disorder](#)
- [Weight Gain](#)