


CASP7 polyclonal antibody

Catalog # PAB19506 Size 100 ug

Applications



Western Blot (Cell lysate)

Western blot analysis of whole cell extracts with CASP7 polyclonal antibody (Cat # PAB19506).

Lane 1, HeLa

Lane 2, MCF-7

Lane 3, SMMC

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of CASP7.
Immunogen	A synthetic peptide corresponding to amino acids at middle region of human CASP7.
Host	Rabbit
Reactivity	Human, Mouse, Rat
Specificity	It different from the related mouse and rat sequence by two amino acids.
Form	Lyophilized
Purification	Immunoaffinity purification
Isotype	IgG
Recommend Usage	Western Blot (1 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ (5 mg BSA, 0.05 mg sodium azide, 0.05 mg Thimerosal)

Storage Instruction

Store at -20°C on dry atmosphere.

After reconstitution with 200 uL of deionized water and concentration will be 500 ug/mL, store at -20°C or lower.

Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide and thimerosal: POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of whole cell extracts with CASP7 polyclonal antibody (Cat # PAB19506).

Lane 1, HeLa

Lane 2, MCF-7

Lane 3, SMMC

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Gene Info — CASP7

Entrez GeneID[840](#)**Gene Name**

CASP7

Gene Alias

CMH-1, ICE-LAP3, MCH3

Gene Description

caspase 7, apoptosis-related cysteine peptidase

Omim ID[601761](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. The precursor of this caspase is cleaved by caspase 3 and 10. It is activated upon cell death stimuli and induces apoptosis. Alternative splicing results in four transcript variants, encoding three distinct isoforms. [provided by RefSeq]

Other Designations

ICE-like apoptotic protease 3|Lice2 alpha/beta/gamma|OTTHUMP00000020510|OTTHUMP00000020511|OTTHUMP00000020512|OTTHUMP00000020514|apoptotic protease MCH-3|caspase 7|caspase 7 isoform delta|caspase 7, apoptosis-related cysteine protease

Pathway

- [Apoptosis](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer Disease](#)
- [Amnesia](#)
- [Arthritis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cognition Disorders](#)
- [Colorectal Neoplasms](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Endometrial Neoplasms](#)
- [Esophageal Neoplasms](#)
- [Gastrointestinal Stromal Tumors](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)

- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Multiple Sclerosis](#)
- [Neoplasm Metastasis](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Neovascularization](#)
- [Neuropsychological Tests](#)
- [Occupational Diseases](#)
- [Prostatic Neoplasms](#)
- [Small Cell Lung Carcinoma](#)
- [Stomach Neoplasms](#)
- [Tobacco Use Disorder](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)