


ANGPTL4 polyclonal antibody

Catalog # PAB19515 Size 100 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded human breast cancer tissue section with ANGPTL4 polyclonal antibody (Cat # PAB19515).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of ANGPTL4.
Immunogen	A synthetic peptide corresponding to amino acids at middle region of human ANGPTL4.
Host	Rabbit
Reactivity	Human
Form	Lyophilized
Purification	Immunoaffinity purification
Isotype	IgG
Recommend Usage	Western Blot (1 ug/mL) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	Lyophilized from 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ (5 mg BSA, 0.05 mg sodium azide, 0.05 mg Thimerosal)

Storage Instruction

Store at -20°C on dry atmosphere.

After reconstitution with 200 uL of deionized water and concentration will be 500 ug/mL, store at -20°C or lower.

Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide and thimerosal: POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded human breast cancer tissue section with ANGPTL4 polyclonal antibody (Cat # PAB19515).

Gene Info — ANGPTL4

Entrez GeneID[51129](#)**Gene Name**

ANGPTL4

Gene Alias

ANGPTL2, ARP4, FIAF, HFARP, NL2, PGAR, pp1158

Gene Description

angiopoietin-like 4

Omim ID[605910](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

This gene is a member of the angiopoietin/angiopoietin-like gene family and encodes a glycosylated, secreted protein with a fibrinogen C-terminal domain. This gene is induced under hypoxic conditions in endothelial cells and is the target of peroxisome proliferation activators. The encoded protein is a serum hormone directly involved in regulating glucose homeostasis, lipid metabolism, and insulin sensitivity and also acts as an apoptosis survival factor for vascular endothelial cells. The encoded protein may play a role in several cancers and it also has been shown to prevent the metastatic process by inhibiting vascular activity as well as tumor cell motility and invasiveness. Decreased expression of this protein has been associated with type 2 diabetes. Alternatively spliced transcript variants encoding different isoforms have been described. This gene was previously referred to as ANGPTL2 but has been renamed ANGPTL4. [provided by RefSeq]

Other Designations

PPARG angiopoietin related protein|angiopoietin-like 4 protein|angiopoietin-related protein 4|fast insulin-induced adipose factor|hepatic angiopoietin-related protein|hepatic fibrinogen/angiopoietin-related protein|peroxisome proliferator-activated receptor (P

Pathway

- [PPAR signaling pathway](#)

Disease

- [Amyotrophic lateral sclerosis](#)
- [Anoxia](#)
- [Atherosclerosis](#)
- [Cardiovascular Diseases](#)
- [Coronary Artery Disease](#)
- [Coronary Disease](#)
- [Diabetes Mellitus](#)
- [Dyslipidemias](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Kidney Failure](#)
- [Syndrome](#)
- [Weight Gain](#)