


EPHB6 polyclonal antibody


Catalog # PAB2074 Size 400 uL

Applications



Western Blot (Cell lysate)


Western blot analysis of EPHB6 polyclonal antibody (Cat # PAB2074) in A-549 cell lysate. EPHB6 (arrow) was detected using purified Polyclonal antibody. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Formalin-fixed and paraffin-embedded human cancer tissue reacted with EPHB6 polyclonal antibody (Cat # PAB2074), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.

This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma.



Flow Cytometry

Flow cytometric analysis of MCF-7 cells using EPHB6 polyclonal antibody (Cat # PAB2074)(bottom histogram) compared to a negative control cell (top histogram).

FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Specification

Product Description

Rabbit polyclonal antibody raised against synthetic peptide of EPHB6.



Product Information

Immunogen	A synthetic peptide (conjugated with KLH) corresponding to N-terminus of human EPHB6.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Protein G purification
Recommend Usage	Western Blot (1:1000) Immunohistochemistry (1:50-100) Flow cytometry (1:10-50) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot (Cell lysate)

Western blot analysis of EPHB6 polyclonal antibody (Cat # PAB2074) in A-549 cell lysate. EPHB6 (arrow) was detected using purified Polyclonal antibody.

Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed and paraffin-embedded human cancer tissue reacted with EPHB6 polyclonal antibody (Cat # PAB2074), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining.

This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma.

Flow Cytometry

Flow cytometric analysis of MCF-7 cells using EPHB6 polyclonal antibody (Cat # PAB2074)(bottom histogram) compared to a negative control cell (top histogram).

FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Gene Info — EPHB6

Entrez GenelD 2051



Product Information

Protein Accession#	NP_004436;O15197
Gene Name	EPHB6
Gene Alias	HEP, MGC129910, MGC129911
Gene Description	EPH receptor B6
Omim ID	602757
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, par ticularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosp hatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The E ph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The ephrin receptor encoded by this gene lacks the kinase activity of most receptor tyrosine kinases and binds to ephrin-B ligands. [provided by RefSeq
Other Designations	ephrin receptor EphB6

Publication Reference

 The EphB6 receptor inhibits JNK activation in T lymphocytes and modulates T cell receptor-mediated responses.

Freywald A, Sharfe N, Rashotte C, Grunberger T, Roifman CM.

The Journal of Biological Chemistry 2003 Mar; 278(12):10150.

Application: WB-Tr, Human, Jurkat cells

• EphB6 crosslinking results in costimulation of T cells.

Luo H, Yu G, Wu Y, Wu J.

The Journal of Clinical Investigation 2002 Oct; 110(8):1141.

Application: Flow Cyt, IF, Human, Human T cells

Multiple roles of EPH receptors and ephrins in neural development.

Wilkinson DG.

Nature Reviews. Neuroscience 2001 Mar; 2(3):155.



Pathway

• Axon guidance