# TBC1D15 polyclonal antibody

Catalog # PAB20818 Size 100 uL

## Applications



#### Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining of human pancreas with TBC1D15 polyclonal antibody (Cat # PAB20818) shows strong cytoplasmic positivity in exocrine glandular cells and islet cells at 1:10-1:20 dilution.

Specification	
Product Description	Rabbit polyclonal antibody raised against recombinant TBC1D15.
Immunogen	Recombinant protein corresponding to amino acids of human TBC1D15.
Sequence	LRVLEKDAEVIVDWRPLDDALDSSSILYARKDSSSVVEWTQAPKERGHRGSEHLNSYEAEWDM VNTVSFKRKPHTNGDAPSHRNGKSKWSFLFSLTDLKSIKQNKEGMGWSYLVFCLKDDVVLPALH F
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
lsotype	lgG
Recommend Usage	Immunohistochemistry (1:10-1:20) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide)

Copyright © 2023 Abnova Corporation. All Rights Reserved.

# 😵 Abnova

### **Product Information**

**Storage Instruction** 

Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.

Allquot to avoid repeated freezing and thavi

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

## Applications

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human pancreas with TBC1D15 polyclonal antibody (Cat # PAB20818) shows strong cytoplasmic positivity in exocrine glandular cells and islet cells at 1:10-1:20 dilution.

Gene Info — TBC1D15	
Entrez GenelD	<u>64786</u>
Protein Accession#	<u>Q8TC07</u>
Gene Name	TBC1D15
Gene Alias	DKFZp686M1379, DKFZp761D0223, FLJ12085
Gene Description	TBC1 domain family, member 15
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a member of the Ras-like proteins in brain-GTPase activating protein superfa mily that share a conserved Tre-2/Bub2/Cdc16 domain. The encoded protein interacts with Ras-li ke protein in brain 5A and may function as a regulator of intracellular trafficking. Alternate splicing results in multiple transcript variants. [provided by RefSeq
Other Designations	-