


SDCCAG8 polyclonal antibody

Catalog # PAB24689

Size 100 ug

Applications



Western Blot (Cell lysate)

SDCCAG8 polyclonal antibody (Cat # PAB24689) (0.3 ug/mL) staining of HepG2 lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Specification

Product Description	Goat polyclonal antibody raised against synthetic peptide of SDCCAG8.
Immunogen	A synthetic peptide corresponding to internal region of human SDCCAG8.
Sequence	CTINDQSQYIHHLEAE
Host	Goat
Theoretical MW (kDa)	80
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Concentration	0.5 mg/mL
Recommend Usage	ELISA (1:128000) Western Blot (0.3-1 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In Tris saline, pH 7.3 (0.5 % BSA, 0.02% sodium azide)

Storage Instruction

Store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

SDCCAG8 polyclonal antibody (Cat # PAB24689) (0.3 ug/mL) staining of HepG2 lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — SDCCAG8

Entrez GeneID[10806](#)**Protein Accession#**[NP_006633.1](#)**Gene Name**

SDCCAG8

Gene Alias

CCCAP, HSPC085, NY-CO-8

Gene Description

serologically defined colon cancer antigen 8

Gene Ontology[Hyperlink](#)**Other Designations**

OTTHUMP00000037913|centrosomal colon cancer autoantigen

Disease

- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [HIV Infections](#)
- [Multiple Sclerosis](#)

- [Obesity](#)
- [Tobacco Use Disorder](#)