


ADORA2A polyclonal antibody


Catalog # PAB24997 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis of HepG2 cell extracts with ADORA2A polyclonal antibody (Cat # PAB24997).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded human skeletal muscle tissue with ADORA2A polyclonal antibody (Cat # PAB24997).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of ADORA2A.
Immunogen	A synthetic peptide corresponding to amino acids 330-380 of human ADORA2A.
Host	Rabbit
Theoretical MW (kDa)	45
Reactivity	Human
Specificity	ADORA2A polyclonal antibody detects endogenous levels of ADORA2A protein.
Form	Liquid

Purification	Antigen affinity purification
Concentration	1 mg/mL
Recommend Usage	Western Blot (1:500-1:1000) Immunohistochemistry (1:50-1:200) Immunofluorescence (1:50-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (0.05% sodium azide).
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of HepG2 cell extracts with ADORA2A polyclonal antibody (Cat # PAB24997).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of paraffin-embedded human skeletal muscle tissue with ADORA2A polyclonal antibody (Cat # PAB24997).

- Immunofluorescence

Gene Info — ADORA2A

Entrez GeneID	135
Protein Accession#	P29274
Gene Name	ADORA2A
Gene Alias	ADORA2, RDC8, hA2aR
Gene Description	adenosine A2a receptor
Omim ID	102776
Gene Ontology	Hyperlink

Gene Summary

This gene encodes a protein which is one of several receptor subtypes for adenosine. The activity of the encoded protein, a G-protein coupled receptor family member, is mediated by G proteins which activate adenylyl cyclase. The encoded protein is abundant in basal ganglia, vasculature and platelets and it is a major target of caffeine. [provided by RefSeq]

Other Designations

adenosine A2 receptor|adenosine receptor subtype A2a

Pathway

- [Calcium signaling pathway](#)
- [Neuroactive ligand-receptor interaction](#)
- [Vascular smooth muscle contraction](#)

Disease

- [Amphetamine-Related Disorders](#)
- [Anorexia Nervosa](#)
- [Anxiety Disorders](#)
- [Arousal](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Autistic Disorder](#)
- [Bulimia](#)
- [Calcinosis](#)
- [Cardiovascular Diseases](#)
- [Cognition](#)
- [Coronary Artery Disease](#)
- [Depressive Disorder](#)
- [Diabetes Mellitus](#)

- [Drug Hypersensitivity](#)
- [Drug Toxicity](#)
- [Edema](#)
- [Fatigue](#)
- [Genetic Predisposition to Disease](#)
- [Headache](#)
- [Huntington disease](#)
- [Hypertension](#)
- [Mental Disorders](#)
- [Migraine with Aura](#)
- [Mood Disorders](#)
- [Neuropsychological Tests](#)
- [Panic Disorder](#)
- [Parkinson disease](#)
- [Personality Inventory](#)
- [Psychiatric Status Rating Scales](#)
- [Psychomotor Performance](#)
- [Psychoses](#)
- [Reaction Time](#)
- [Schizophrenia](#)
- [Schizophrenic Psychology](#)
- [Sleep](#)
- [Sleep Disorders](#)
- [Sleep Initiation and Maintenance Disorders](#)
- [Substance Withdrawal Syndrome](#)
- [Substance-Related Disorders](#)

- [Syncope](#)
- [Weight Gain](#)