


ATF4 (phospho S245) polyclonal antibody

Catalog # PAB25032 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue with ATF4 (phospho S245) polyclonal antibody (Cat # PAB25032).

Specification	
Product Description	Rabbit polyclonal antibody raised against synthetic phosphopeptide of ATF4.
Immunogen	Synthetic phosphopeptide corresponding to residues surrounding S245 of ATF4.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Concentration	1 mg/mL
Purity	> 95% by SDS-PAGE
Recommend Usage	Western Blot (1:500-1:1000) Immunohistochemistry (1:50-1:200) Immunoprecipitation (1:50-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (0.05% sodium azide)



Product Information

Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)
 Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue with ATF4 (phospho S245) polyclonal antibody (Cat # PAB25032).
- Immunoprecipitation

Gene Info — ATF4	
Entrez GenelD	<u>468</u>
Gene Name	ATF4
Gene Alias	CREB-2, CREB2, TAXREB67, TXREB
Gene Description	activating transcription factor 4 (tax-responsive enhancer element B67)
Omim ID	<u>604064</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a transcription factor that was originally identified as a widely expressed mam malian DNA binding protein that could bind a tax-responsive enhancer element in the LTR of HTL V-1. The encoded protein was also isolated and characterized as the cAMP-response element binding protein 2 (CREB-2). The protein encoded by this gene belongs to a family of DNA-binding proteins that includes the AP-1 family of transcription factors, cAMP-response element binding proteins (CREBs) and CREB-like proteins. These transcription factors share a leucine zipper region that is involved in protein-protein interactions, located C-terminal to a stretch of basic amino acids that functions as a DNA binding domain. Two alternative transcripts encoding the same protein have been described. Two pseudogenes are located on the X chromsome at q28 in a region containing a large inverted duplication. [provided by RefSeq
Other Designations	activating transcription factor 4 cAMP response element-binding protein 2



Pathway

- GnRH signaling pathway
- Long-term potentiation
- MAPK signaling pathway
- Neurotrophin signaling pathway
- Prostate cancer

Disease

- Bipolar Disorder
- Genetic Predisposition to Disease
- Mental Disorders
- Neuropsychological Tests
- Schizophrenia