


CCKAR polyclonal antibody

Catalog # PAB25584

Size 50 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of formalin-fixed, paraffin-embedded human prostate (A) and human adrenal gland (B) tissue after heat-induced antigen retrieval.

Using CCKAR polyclonal antibody (Cat # PAB25584).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of CCKAR.
Immunogen	A synthetic peptide corresponding to 20 amino acids at internal region of human CCKAR.
Host	Rabbit
Reactivity	Human
Specificity	BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Form	Liquid
Purification	Immunoaffinity chromatography
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (10 ug/mL) The optimal working dilution should be determined by the end user.

Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of formalin-fixed, paraffin-embedded human prostate (A) and human adrenal gland (B) tissue after heat-induced antigen retrieval.

Using CCKAR polyclonal antibody (Cat # PAB25584).

Gene Info — CCKAR

Entrez GeneID	886
Gene Name	CCKAR
Gene Alias	CCK-A, CCK1-R, CCKRA
Gene Description	cholecystokinin A receptor
Omim ID	118444
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a G-protein coupled receptor that binds non-sulfated members of the cholecystokinin (CCK) family of peptide hormones. This receptor is a major physiologic mediator of pancreatic enzyme secretion and smooth muscle contraction of the gallbladder and stomach. In the central and peripheral nervous system this receptor regulates satiety and the release of beta-endorphin and dopamine. [provided by RefSeq]
Other Designations	cholecystokinin type-A receptor cholecystokinin-1 receptor

Pathway

- [Calcium signaling pathway](#)
- [Neuroactive ligand-receptor interaction](#)

Disease

- [Alcohol Withdrawal Delirium](#)
- [Alcoholism](#)
- [Alcohol-Related Disorders](#)
- [Anorexia Nervosa](#)
- [Bipolar Disorder](#)
- [Bulimia](#)
- [Bulimia Nervosa](#)
- [Cardiovascular Diseases](#)
- [Depressive Disorder](#)
- [Diabetes Mellitus](#)
- [Disease Models](#)
- [Drug Toxicity](#)
- [Eating Disorders](#)
- [Edema](#)
- [Gallbladder Neoplasms](#)
- [Gallstones](#)
- [Genetic Predisposition to Disease](#)
- [Hallucinations](#)
- [Hypercholesterolemia](#)
- [Hyperparathyroidism](#)
- [Insomnia](#)
- [Liver Diseases](#)
- [Mental Disorders](#)
- [Obesity](#)

- [Panic Disorder](#)
- [Parkinson disease](#)
- [Psychiatric Status Rating Scales](#)
- [Schizophrenia](#)
- [Schizophrenic Psychology](#)
- [Sleep Apnea](#)
- [Tobacco Use Disorder](#)
- [Wechsler Scales](#)
- [Weight Gain](#)