


CASP3 polyclonal antibody

Catalog # PAB25662 Size 100 ug

Applications



Immunohistochemistry

Immunohistochemical analysis of CASP3 in human tonsil tissue with CASP3 polyclonal antibody (Cat # PAB25662) at 5 ug/mL.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of CASP3.
Immunogen	A synthetic peptide corresponding to 17 amino acids at internal region of human CASP3.
Host	Rabbit
Theoretical MW (kDa)	30
Reactivity	Human
Specificity	At least two isoforms of Caspase-3 are known to exist.
Form	Liquid
Purification	Peptide affinity purification
Concentration	1 mg/mL
Isotype	IgG
Recommend Usage	Immunohistochemistry (5 ug/mL) The optimal working dilution should be determined by the end user.

Storage Buffer	In PBS (0.05% sodium azide)
Storage Instruction	Store at 4°C for three months. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry

Immunohistochemical analysis of CASP3 in human tonsil tissue with CASP3 polyclonal antibody (Cat # PAB25662) at 5 ug/mL.

- Enzyme-linked Immunoabsorbent Assay

Gene Info — CASP3

Entrez GeneID	836
Protein Accession#	NP_004337
Gene Name	CASP3
Gene Alias	CPP32, CPP32B, SCA-1
Gene Description	caspase 3, apoptosis-related cysteine peptidase
Omim ID	600636
Gene Ontology	Hyperlink

Gene Summary	This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 6, 7 and 9, and the protein itself is processed by caspases 8, 9 and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. Alternative splicing of this gene results in two transcript variants that encode the same protein. [provided by RefSeq]
--------------	---

Other Designations	OTTHUMP00000165054 PARP cleavage protease SREBP cleavage activity 1 Yama apopain caspase 3 caspase 3, apoptosis-related cysteine protease cysteine protease CPP32 procaspase3
--------------------	---

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Colorectal cancer](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [MAPK signaling pathway](#)
- [Natural killer cell mediated cytotoxicity](#)
- [p53 signaling pathway](#)
- [Pathways in cancer](#)

Disease

- [Adenocarcinoma](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Clubfoot](#)
- [Colorectal Neoplasms](#)
- [Diabetes Mellitus](#)
- [Diabetic Nephropathies](#)
- [Disease Progression](#)
- [Edema](#)
- [Endometrial Neoplasms](#)
- [Esophageal Neoplasms](#)

- [Gastrointestinal Stromal Tumors](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Lung carcinoma](#)
- [Lung Neoplasms](#)
- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Myeloma](#)
- [Multiple Sclerosis](#)
- [NARP](#)
- [Neoplasm Metastasis](#)
- [Neoplasms](#)
- [Neovascularization](#)
- [Occupational Diseases](#)
- [Prostatic Neoplasms](#)
- [Pulmonary Disease](#)
- [Small Cell Lung Carcinoma](#)
- [Stomach Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Waldenstrom Macroglobulinemia](#)

- [Werner syndrome](#)