


ATF2 polyclonal antibody


Catalog # PAB25947 Size 100 ug

Applications



Western Blot (Cell lysate)

Western blot analysis of extracts from MDA-MB-435 cells using ATF2 polyclonal antibody (Cat # PAB25947).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using ATF2 polyclonal antibody (Cat # PAB25947).

Specification	
Product Description	Rabbit polyclonal antibody raised against synthetic peptide of ATF2.
Immunogen	A synthetic peptide corresponding to residues surrounding S112/S94 of human ATF2.
Sequence	D-L-Sp-P-L
Host	Rabbit
Reactivity	Human, Mouse, Rat
Form	Liquid



Product Information

Purification	Affinity chromatography
Concentration	1 mg/mL
Recommend Usage	Immunohistochemistry (1:50-1:100)
	Western Blot (1:500-1:1000)
	The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20°C.
	Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul
	d be handled by trained staff only.

Applications

Western Blot (Cell lysate)

Western blot analysis of extracts from MDA-MB-435 cells using ATF2 polyclonal antibody (Cat # PAB25947).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using ATF2 polyclonal antibody (Cat # PAB25947).

Gene Info — ATF2	
Entrez GeneID	1386
Protein Accession#	P15336
Gene Name	ATF2
Gene Alias	CRE-BP1, CREB2, HB16, MGC111558, TREB7
Gene Description	activating transcription factor 2
Omim ID	123811
Gene Ontology	<u>Hyperlink</u>



Product Information

Gene Summary

This gene encodes a transcription factor that is a member of the leucine zipper family of DNA bin ding proteins. This protein binds to the cAMP-responsive element (CRE), an octameric palindrom e. The protein forms a homodimer or heterodimer with c-Jun and stimulates CRE-dependent tran scription. The protein is also a histone acetyltransferase (HAT) that specifically acetylates histone s H2B and H4 in vitro; thus it may represent a class of sequence-specific factors that activate tran scription by direct effects on chromatin components. Additional transcript variants have been iden tified but their biological validity has not been determined. [provided by RefSeq

Other Designations

OTTHUMP00000163262|activating transcription factor 2 splice variant ATF2-var2|cAMP respons ive element binding protein 2, formerly

Pathway

MAPK signaling pathway

Disease

- Bipolar Disorder
- Cardiovascular Diseases
- Diabetes Mellitus
- Edema
- Genetic Predisposition to Disease
- Schizophrenia
- Tobacco Use Disorder