


CCR6 polyclonal antibody

Catalog # PAB26134

Size 50 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human tonsil with CCR6 polyclonal antibody (Cat # PAB26134).

Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of CCR6.
Immunogen	A synthetic peptide corresponding to 20 amino acids at C-terminus of human CCR6.
Host	Rabbit
Reactivity	Human, Monkey
Specificity	BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Form	Liquid
Purification	Immunoaffinity chromatography
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (5 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human tonsil with CCR6 polyclonal antibody (Cat # PAB26134).

Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Gene Info — CCR6

Entrez GeneID [1235](#)

Protein Accession# [P51684](#)

Gene Name CCR6

Gene Alias BN-1, CD196, CKR-L3, CKR6, CKRL3, CMKBR6, DCR2, DRY-6, GPR-CY4, GPR29, GPRCY4, STRL22

Gene Description chemokine (C-C motif) receptor 6

Omim ID [601835](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. The gene is preferentially expressed by immature dendritic cells and memory T cells. The ligand of this receptor is macrophage inflammatory protein 3 alpha (MIP-3 alpha). This receptor has been shown to be important for B-lineage maturation and antigen-driven B-cell differentiation, and it may regulate the migration and recruitment of dendritic and T cells during inflammatory and immunological responses. Alternatively spliced transcript variants that encode the same protein have been described for this gene. [provided by RefSeq]

Other Designations G protein-coupled receptor 29|OTTHUMP00000017618|OTTHUMP00000017619|chemokine (C-C) receptor 6|chemokine receptor-like 3|seven-transmembrane receptor, lymphocyte, 22

Pathway

- [Chemokine signaling pathway](#)
- [Cytokine-cytokine receptor interaction](#)

Disease

- [Acquired Immunodeficiency Syndrome](#)
- [Arthritis](#)
- [Birth Weight](#)
- [Cleft Lip](#)
- [Cleft Palate](#)
- [Crohn Disease](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [HIV Infections](#)
- [Inflammatory Bowel Diseases](#)
- [Leukemia](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Tobacco Use Disorder](#)
- [Vitiligo](#)