


TACR1 polyclonal antibody

Catalog # PAB26420

Size 50 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human brain, neurons and glia with TACR1 polyclonal antibody (Cat # PAB26420). Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of TACR1.
Immunogen	A synthetic peptide corresponding to 16 amino acids at 2nd extracellular domain of human TACR1.
Host	Rabbit
Reactivity	Bovine, Dog, Human, Monkey
Specificity	BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Form	Liquid
Purification	Immunoaffinity chromatography
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (2-4 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human brain, neurons and glia with TACR1 polyclonal antibody (Cat # PAB26420). Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Gene Info — TACR1

Entrez GeneID [6869](#)

Protein Accession# [P25103](#)

Gene Name TACR1

Gene Alias NK1R, NK1R, SPR, TAC1R

Gene Description tachykinin receptor 1

Omim ID [162323](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene belongs to a gene family of tachykinin receptors. These tachykinin receptors are characterized by interactions with G proteins and contain seven hydrophobic transmembrane regions. This gene encodes the receptor for the tachykinin substance P, also referred to as neurokinin 1. The encoded protein is also involved in the mediation of phosphatidylinositol metabolism of substance P. [provided by RefSeq]

Other Designations NK-1 receptor|neurokinin 1 receptor|tachykinin 1 receptor (substance P receptor, neurokinin 1 receptor)|tachykinin receptor 1 (substance P receptor; neurokinin-1 receptor)

Pathway

- [Calcium signaling pathway](#)
- [Neuroactive ligand-receptor interaction](#)

Disease

- [Alcoholism](#)
- [Anorexia Nervosa](#)
- [Asthma](#)
- [Bipolar Disorder](#)
- [Bulimia](#)
- [Chronic Disease](#)
- [Constipation](#)
- [Cough](#)
- [Endometriosis](#)
- [Fibromyalgia](#)
- [Genetic Predisposition to Disease](#)
- [Mental Disorders](#)
- [Nausea](#)
- [Pain](#)
- [Panic Disorder](#)
- [Pregnancy Complications](#)
- [Radius Fractures](#)
- [Recurrence](#)
- [Reflex Sympathetic Dystrophy](#)
- [Schizophrenia](#)
- [Tobacco Use Disorder](#)