


GPR85 polyclonal antibody

Catalog # PAB26480 Size 50 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human putamen, neurons with GPR85 polyclonal antibody (Cat # PAB26480). Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Specification	
Product Description	Rabbit polyclonal antibody raised against synthetic peptide of GPR85.
Immunogen	A synthetic peptide corresponding to 18 amino acids at 3rd cytoplasmic domain of human GPR85.
Host	Rabbit
Reactivity	Bovine, Chicken, Dog, Hamster, Horse, Human, Monkey, Mouse, Pig, Rabbit, Rat
Specificity	BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Form	Liquid
Purification	Immunoaffinity chromatography
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (10-19 ug/mL) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.



Product Information

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical (Formalin/PFA-fixed paraffin-embedded sections) staining in human putamen, neurons with GPR85 polyclonal antibody (Cat # PAB26480). Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Gene Info — GPR85	
Entrez GenelD	<u>54329</u>
Protein Accession#	P60893
Gene Name	GPR85
Gene Alias	SREB, SREB2
Gene Description	G protein-coupled receptor 85
Omim ID	605188
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Members of the G protein-coupled receptor (GPCR) family, such as GPR85, have a similar struct ure characterized by 7 transmembrane domains. Activation of GPCRs by extracellular stimuli, such as neurotransmitters, hormones, or light, induces an intracellular signaling cascade mediated by heterotrimeric GTP-binding proteins, or G proteins (Matsumoto et al., 2000 [PubMed 10833454]). [supplied by OMIM
Other Designations	seven transmembrane helix receptor super conserved receptor expressed in brain 2