


BCL2 polyclonal antibody


Catalog # PAB26647 Size 100 ug

Applications



Western Blot (Cell lysate)

Western blot analysis of extracts from MCF-7 cells using BCL2 polyclonal antibody (Cat # PAB26647).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using BCL2 polyclonal antibody (Cat # PAB26647).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of BCL2.
Immunogen	A synthetic peptide corresponding to residues surrounding S70 of human BCL2.
Sequence	R-T-Sp-P-L
Host	Rabbit
Theoretical MW (kDa)	26
Reactivity	Human

Form	Liquid
Purification	Affinity chromatography
Concentration	1 mg/mL
Recommend Usage	Western Blot (1:500-1:1000) Immunohistochemistry (1:50-1:100) Immunofluorescence (1:100-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (without Mg ²⁺ and Ca ²⁺), 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot (Cell lysate)

Western blot analysis of extracts from MCF-7 cells using BCL2 polyclonal antibody (Cat # PAB26647).

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using BCL2 polyclonal antibody (Cat # PAB26647).

- Immunofluorescence

Gene Info — BCL2

Entrez GeneID	596
Protein Accession#	P10415
Gene Name	BCL2
Gene Alias	Bcl-2
Gene Description	B-cell CLL/lymphoma 2
Omim ID	151430
Gene Ontology	Hyperlink

Gene Summary

This gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends. [provided by RefSeq]

Other Designations

B-cell lymphoma protein 2|OTTHUMP00000163680

Pathway

- [Amyotrophic lateral sclerosis \(ALS\)](#)
- [Apoptosis](#)
- [Colorectal cancer](#)
- [Focal adhesion](#)
- [Neurotrophin signaling pathway](#)
- [Pathways in cancer](#)
- [Prostate cancer](#)
- [Small cell lung cancer](#)

Disease

- [Adenocarcinoma](#)
- [Alzheimer Disease](#)
- [Amnesia](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Azoospermia](#)
- [Bile Duct Neoplasms](#)
- [Brain Injuries](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)

- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cholangiocarcinoma](#)
- [Chromosome Aberrations](#)
- [Clubfoot](#)
- [Cognition](#)
- [Cognition Disorders](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Head and Neck Neoplasms](#)
- [Hematologic Diseases](#)
- [Hodgkin Disease](#)
- [Kidney Failure](#)
- [Kidney Neoplasms](#)
- [Laryngeal Neoplasms](#)
- [Leukemia](#)
- [Leukemia-Lymphoma](#)
- [Lung Neoplasms](#)
- [Lupus Erythematosus](#)
- [Lymphatic Metastasis](#)
- [Lymphoma](#)
- [Lymphoproliferative Disorders](#)

- [Multiple Sclerosis](#)
- [NARP](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Recurrence](#)
- [Neovascularization](#)
- [Neuropsychological Tests](#)
- [Occupational Diseases](#)
- [Oropharyngeal Neoplasms](#)
- [Ovarian Neoplasms](#)
- [Postoperative Complications](#)
- [Prostate cancer](#)
- [Prostatic Neoplasms](#)
- [Prosthesis Failure](#)
- [Pulmonary Disease](#)
- [Thyroid Neoplasms](#)
- [Translocation](#)
- [Waldenstrom Macroglobulinemia](#)
- [Werner syndrome](#)