


CAMK2A polyclonal antibody

Catalog # PAB26799 Size 100 ug

Applications



Western Blot (Tissue lysate)

Western blot analysis of extracts from mouse brain tissue using CAMK2A polyclonal antibody (Cat # PAB26799).

| Specification | |
|---------------------|---|
| Product Description | Rabbit polyclonal antibody raised against synthetic peptide of CAMK2A. |
| Immunogen | A synthetic peptide corresponding to residues surrounding T286 of human CAMK2A. |
| Sequence | Q-E-Tp-V-D |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Form | Liquid |
| Purification | Affinity chromatography |
| Concentration | 1 mg/mL |
| Recommend Usage | Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user. |
| Storage Buffer | In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide) |



Product Information

| Storage Instruction | Store at -20°C. Aliquot to avoid repeated freezing and thawing. |
|---------------------|---|
| Note | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only. |

Applications

• Western Blot (Tissue lysate)

Western blot analysis of extracts from mouse brain tissue using CAMK2A polyclonal antibody (Cat # PAB26799).

| Gene Info — CAMK2A | |
|--------------------|--|
| Entrez GenelD | <u>815</u> |
| Protein Accession# | Q9UQM7 |
| Gene Name | CAMK2A |
| Gene Alias | CAMKA, KIAA0968 |
| Gene Description | calcium/calmodulin-dependent protein kinase II alpha |
| Omim ID | 114078 |
| Gene Ontology | Hyperlink |
| Gene Summary | The product of this gene belongs to the serine/threonine protein kinases family, and to the Ca(2+)/calmodulin-dependent protein kinases subfamily. Calcium signaling is crucial for several aspects of plasticity at glutamatergic synapses. This calcium calmodulin-dependent protein kinase is composed of four different chains: alpha, beta, gamma, and delta. The alpha chain encoded by this gene is required for hippocampal long-term potentiation (LTP) and spatial learning. In addition to its calcium-calmodulin (CaM)-dependent activity, this protein can undergo autophosphorylation, resulting in CaM-independent activity. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq |
| Other Designations | CaM kinase II alpha subunit CaM-kinase II alpha chain CaMK-II alpha subunit CaMKIINalpha OTT HUMP00000165787 OTTHUMP00000165788 calcium/calmodulin-dependent protein kinase (Ca M kinase) II alpha calcium/calmodulin-dependent protein kinase II alpha-B subunit |

Pathway

Calcium signaling pathway



- ErbB signaling pathway
- Glioma
- GnRH signaling pathway
- Long-term potentiation
- Melanogenesis
- Neurotrophin signaling pathway
- Olfactory transduction
- Wnt signaling pathway

Disease

- Bipolar Disorder
- Cognition
- Genetic Predisposition to Disease
- Schizophrenia
- Schizophrenic Psychology
- Tobacco Use Disorder
- Weight Gain