


CCNB1 polyclonal antibody

Catalog # PAB26851 Size 100 ug


Applications



Peptide - +

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using CCNB1 polyclonal antibody (Cat # PAB26851).



Immunofluorescence

Immunofluorescence staining of methanol-fixed HeLa cells using CCNB1 polyclonal antibody (Cat # PAB26851).

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of CCNB1.
Immunogen	A synthetic peptide (conjugated with KLH) corresponding to residues surrounding S147 of human CCNB1.
Sequence	A-F-Sp-D-V
Host	Rabbit
Theoretical MW (kDa)	60
Reactivity	Human

Form	Liquid
Purification	Affinity chromatography
Concentration	1 mg/mL
Recommend Usage	Immunohistochemistry (1:50-1:100) Immunofluorescence (1:100-1:200) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using CCNB1 polyclonal antibody (Cat # PAB26851).

- Immunofluorescence

Immunofluorescence staining of methanol-fixed HeLa cells using CCNB1 polyclonal antibody (Cat # PAB26851).

Gene Info — CCNB1

Entrez GeneID	891
Protein Accession#	P14635
Gene Name	CCNB1
Gene Alias	CCNB
Gene Description	cyclin B1
Omim ID	123836
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a regulatory protein involved in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). Two alternative transcripts have been found, a constitutively expressed transcript and a cell cycle-regulated transcript, that is expressed predominantly during G2/M phase. The different transcripts result from the use of alternate transcription initiation sites. [provided by RefSeq]

Other Designations

G2/mitotic-specific cyclin B1

Pathway

- [Cell cycle](#)
- [p53 signaling pathway](#)

Disease

- [Adenocarcinoma](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [Ovarian Neoplasms](#)