


NTRK2 polyclonal antibody

Catalog # PAB28366

Size 50 ug

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) of human brain neurons with NTRK2 polyclonal antibody (Cat # PAB28366). Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of NTRK2.
Immunogen	A synthetic peptide corresponding to 12 amino acids at cytoplasmic domain of human NTRK2.
Host	Rabbit
Reactivity	Horse, Human, Monkey, Rabbit
Specificity	BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Form	Liquid
Purification	Immunoaffinity chromatography
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (20 ug/ml) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) of human brain neurons with NTRK2 polyclonal antibody (Cat # PAB28366). Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Gene Info — NTRK2

Entrez GeneID [4915](#)

Protein Accession# [Q16620](#)

Gene Name NTRK2

Gene Alias GP145-TrkB, TRKB

Gene Description neurotrophic tyrosine kinase, receptor, type 2

Omim ID [600456](#)

Gene Ontology [Hyperlink](#)

Gene Summary This gene encodes a member of the neurotrophic tyrosine receptor kinase (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. Signalling through this kinase leads to cell differentiation. Mutations in this gene have been associated with obesity and mood disorders. Alternate transcriptional splice variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations BDNF/NT-3 growth factors receptor|OTTHUMP00000021573|OTTHUMP00000021574|OTTHUMP00000021575|OTTHUMP00000021576|OTTHUMP00000021577|tyrosine kinase receptor B

Pathway

- [MAPK signaling pathway](#)
- [Neurotrophin signaling pathway](#)

Disease

- [Alcoholism](#)
- [Alzheimer disease](#)
- [Anorexia Nervosa](#)
- [Antisocial Personality Disorder](#)
- [Asperger Syndrome](#)
- [Attention Deficit Disorder with Hyperactivity](#)
- [Autistic Disorder](#)
- [Avoidance Learning](#)
- [Bipolar Disorder](#)
- [Bulimia](#)
- [Cardiovascular Diseases](#)
- [Cognition](#)
- [Depressive Disorder](#)
- [Diabetes Mellitus](#)
- [Disease Models](#)
- [Eating Disorders](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Mental Disorders](#)
- [Mood Disorders](#)
- [Obsessive-Compulsive Disorder](#)
- [Psychiatric Status Rating Scales](#)
- [Schizophrenia](#)
- [Schizophrenic Psychology](#)

- [Sleep Disorders](#)
- [Social Perception](#)
- [Stress Disorders](#)
- [Tobacco Use Disorder](#)
- [Weight Gain](#)