


ASCC2 polyclonal antibody


Catalog # PAB28435 Size 100 uL

Applications




Western Blot

Western blot analysis of Lane 1: RT-4, Lane 2: EFO-21, Lane 3: A-431, Lane 4: Liver, Lane 5: Tonsil with ASCC2 polyclonal antibody (Cat # PAB28435) at 1:250-1:500 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining of human duodenum with ASCC2 polyclonal antibody (Cat # PAB28435) shows strong cytoplasmic positivity in glandular cells at 1:20-1:50 dilution.



Immunofluorescence

Immunofluorescent staining of U-2 OS with ASCC2 polyclonal antibody (Cat # PAB28435) at 1-4 ug/mL dilution shows positivity in nucleus, cytoplasm & focal adhesions.



Product Description	Rabbit polyclonal antibody raised against recombinant ASCC2.
Immunogen	Recombinant protein corresponding to amino acids of recombinant ASCC2.
Sequence	YSVVVEEVPLQPGESLPYHSVYYEDEYDDTYDGNQVGANDADSDDELISRRPFTIPQVLRTKVPR EGQEEDDDDEEDDADEEAPKPDHFVQDPAVLREKAEARRMAFLAKKGYRHDSSTAVAGSPRG HGQSRETTQERRKKEANKATRANH
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Isotype	lgG
Recommend Usage	Immunohistochemistry(1:20-1:50) Western Blot(1:250-1:500) Immunofluorescence (1-4 ug/ml) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot

Western blot analysis of Lane 1: RT-4, Lane 2: EFO-21, Lane 3: A-431, Lane 4: Liver, Lane 5: Tonsil with ASCC2 polyclonal antibody (Cat # PAB28435) at 1:250-1:500 dilution.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human duodenum with ASCC2 polyclonal antibody (Cat # PAB28435) shows strong cytoplasmic positivity in glandular cells at 1:20-1:50 dilution.

Immunofluorescence

Immunofluorescent staining of U-2 OS with ASCC2 polyclonal antibody (Cat # PAB28435) at 1-4 ug/mL dilution shows positivity in nucleus, cytoplasm & focal adhesions.



Gene Info — ASCC2	
Entrez GenelD	<u>84164</u>
Protein Accession#	<u>F5H6J9</u>
Gene Name	ASCC2
Gene Alias	ASC1p100
Gene Description	activating signal cointegrator 1 complex subunit 2
Gene Ontology	<u>Hyperlink</u>
Other Designations	ASC 1 complex subunit P100 ASC-1 complex subunit P100