


STK39 (phospho S325) polyclonal antibody

Catalog # PAB29219 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining of human brain tissue by STK39 (phospho S325) polyclonal antibody (Cat # PAB29219) without blocking peptide (A) or preincubated with blocking peptide (B) under 1:50-1:100 dilution.

Specification	
Product Description	Rabbit polyclonal antibody raised against synthetic phosphopeptide of human STK39.
Immunogen	Synthetic phosphopeptide (conjugated with KLH) corresponding to residues surrounding S325 of hu man STK39.
Host	Rabbit
Theoretical MW (kDa)	59
Reactivity	Human, Mouse, Rat
Specificity	This antibody detects endogenous levels of STK39 only when phosphorylated at serine 325.
Form	Liquid



Product Information

Purification	Affinity purification
Recommend Usage	Immunohistochemistry (1:50-1:100) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (without Mg ²⁺ and Ca ²⁺), (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Avoid repeated freezing and thawing.

Applications

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human brain tissue by STK39 (phospho S325) polyclonal antibody (Cat # PAB29219) without blocking peptide (A) or preincubated with blocking peptide (B) under 1:50-1:100 dilution.

Gene Info — STK39	
Entrez GenelD	<u>27347</u>
Protein Accession#	Q9UEW8
Gene Name	STK39
Gene Alias	DCHT, DKFZp686K05124, PASK, SPAK
Gene Description	serine threonine kinase 39 (STE20/SPS1 homolog, yeast)
Omim ID	607648
Gene Ontology	<u>Hyperlink</u>
Gene Summary	This gene encodes a serine/threonine kinase that is thought to function in the cellular stress response pathway. The kinase is activated in response to hypotonic stress, leading to phosphorylation of several cation-chloride-coupled cotransporters. The catalytically active kinase specifically activates the p38 MAP kinase pathway, and its interaction with p38 decreases upon cellular stress, suggesting that this kinase may serve as an intermediate in the response to cellular stress. [provide d by RefSeq
Other Designations	Ste20-like protein kinase proline-alanine-rich STE20-related kinase small intestine SPAK-like kin ase

Disease



- Autistic Disorder
- Carcinoma
- Genetic Predisposition to Disease
- Hypertension
- Lung Neoplasms
- Tobacco Use Disorder