


FLT1 (phospho Y1048) polyclonal antibody

Catalog # PAB29223 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human brain tissue by FLT1 (phospho Y1048) polyclonal antibody (Cat # PAB29223) without blocking peptide (A) or preincubated with blocking peptide (B) under 1:50-1:100 dilution.

Specification

Product Description	Rabbit polyclonal antibody raised against synthetic phosphopeptide of human FLT1.
Immunogen	Synthetic phosphopeptide (conjugated with KLH) corresponding to residues surrounding Y1048 of human FLT1.
Host	Rabbit
Theoretical MW (kDa)	150
Reactivity	Human, Mouse, Rat
Specificity	This antibody detects endogenous levels of FLT1 only when phosphorylated at tyrosine 1048.
Form	Liquid

Purification	Affinity purification
Recommend Usage	Immunohistochemistry (1:50-1:100) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.4 (without Mg ²⁺ and Ca ²⁺), (50% glycerol, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Avoid repeated freezing and thawing.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining of human brain tissue by FLT1 (phospho Y1048) polyclonal antibody (Cat # PAB29223) without blocking peptide (A) or preincubated with blocking peptide (B) under 1:50-1:100 dilution.

Gene Info — FLT1

Entrez GeneID	2321
Protein Accession#	P17948
Gene Name	FLT1
Gene Alias	FLT, VEGFR1
Gene Description	fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor)
Omim ID	165070
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a member of the vascular endothelial growth factor receptor (VEGFR) family. VEGFR family members are receptor tyrosine kinases (RTKs) which contain an extracellular ligand-binding region with seven immunoglobulin (Ig)-like domains, a transmembrane segment, and a tyrosine kinase (TK) domain within the cytoplasmic domain. This protein binds to VEGFR-A, VEGFR-B and placental growth factor and plays an important role in angiogenesis and vasculogenesis. Expression of this receptor is found in vascular endothelial cells, placental trophoblast cells and peripheral blood monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Isoforms include a full-length transmembrane receptor isoform and shortened, soluble isoforms. The soluble isoforms are associated with the onset of pre-eclampsia
Other Designations	fms-related tyrosine kinase 1 soluble VEGF receptor 1-14 soluble VEGFR1 variant 2 soluble VEGFR1 variant 21 vascular endothelial growth factor/vascular permeability factor receptor

Pathway

- [Cytokine-cytokine receptor interaction](#)
- [Endocytosis](#)
- [Focal adhesion](#)

Disease

- [Abortion](#)
- [Adenocarcinoma](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Cardiovascular Diseases](#)
- [Cell Transformation](#)
- [Chorioamnionitis](#)
- [Colorectal Neoplasms](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Fetal Growth Retardation](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Hypercholesterolemia](#)
- [Hypersensitivity](#)
- [Inflammation](#)
- [Kidney Failure](#)

- [Lymphoma](#)
- [Malaria](#)
- [Melanoma](#)
- [Neovascularization](#)
- [Obstetric Labor](#)
- [Placenta Diseases](#)
- [Pre-Eclampsia](#)
- [Pregnancy Complications](#)
- [Premature Birth](#)
- [Sarcoidosis](#)
- [Scleroderma](#)
- [Skin Neoplasms](#)
- [Vaginosis](#)