


NRG1 polyclonal antibody

Catalog # PAB30439

Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lateral ventricle with NRG1 polyclonal antibody (Cat # PAB30439) shows weak positivity at 1:500-1:1000 dilution.

Specification

Product Description	Rabbit polyclonal antibody raised against partial recombinant human NRG1.
Immunogen	Recombinant protein corresponding to human NRG1.
Sequence	KWFKNGNELNRKNKPQNIQKKPGKSELRINKASLADSGEYMCKVISKLGNDSASANITVESNAT STSTTGTSHLVKAEKEKTFVNGGECFMVKDLSNPSRYLC
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:500-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide).

Storage Instruction

Store at 4°C. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lateral ventricle with NRG1 polyclonal antibody (Cat # PAB30439) shows weak positivity at 1:500-1:1000 dilution.

Gene Info — NRG1

Entrez GeneID
[3084](#)
Protein Accession#
[Q02297](#)
Gene Name

NRG1

Gene Alias

ARIA, GGF, GGF2, HGL, HRG, HRG1, HRGA, NDF, SMDF

Gene Description

neuregulin 1

Omim ID
[142445 603013](#)
Gene Ontology
[Hyperlink](#)
Gene Summary

The protein encoded by this gene was originally identified as a 44-kD glycoprotein that interacts with the NEU/ERBB2 receptor tyrosine kinase to increase its phosphorylation on tyrosine residues. This protein is a signaling protein that mediates cell-cell interactions and plays critical roles in the growth and development of multiple organ systems. It is known that an extraordinary variety of different isoforms are produced from this gene through alternative promoter usage and splicing. These isoforms are tissue-specifically expressed and differ significantly in their structure, and therefore these isoforms are classified into types I, II, III, IV, V and VI. The gene dysregulation has been linked to diseases such as cancer, schizophrenia and bipolar disorder (BPD). [provided by RefSeq]

Other Designations

glial growth factor|heregulin, alpha (45kD, ERBB2 p185-activator)|neu differentiation factor|neuregulin 1 isoform HRG-gamma|sensory and motor neuron derived factor

Pathway

- [ErbB signaling pathway](#)

Disease

- [Alzheimer disease](#)
- [Amphetamine-Related Disorders](#)
- [Anoxia](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Atrophy](#)
- [Attention](#)
- [Autistic Disorder](#)
- [Bipolar Disorder](#)
- [Cardiovascular Diseases](#)
- [Carotid Artery Diseases](#)
- [Chronic Disease](#)
- [Cognition](#)
- [Cognition Disorders](#)
- [Depressive Disorder](#)
- [Diabetes Mellitus](#)
- [Dominance](#)
- [Edema](#)
- [Extraversion \(Psychology\)](#)
- [Genetic Predisposition to Disease](#)
- [Hirschsprung Disease](#)
- [Infant](#)
- [Inhibition \(Psychology\)](#)
- [Intelligence Tests](#)

- [Leukomalacia](#)
- [Memory](#)
- [Mental Disorders](#)
- [Mental Recall](#)
- [Mental Retardation](#)
- [Neuropsychological Tests](#)
- [Neurotic Disorders](#)
- [Obstetric Labor Complications](#)
- [Perceptual Disorders](#)
- [Personality Tests](#)
- [Psychiatric Status Rating Scales](#)
- [Psychomotor Performance](#)
- [Psychoses](#)
- [Psychotic Disorders](#)
- [Reaction Time](#)
- [Recurrence](#)
- [Schizophrenia](#)
- [Schizophrenic Psychology](#)
- [Schizotypal Personality Disorder](#)
- [Space Perception](#)
- [Speech Disorders](#)
- [Startle Reaction](#)
- [Stress](#)
- [Task Performance and Analysis](#)
- [Tobacco Use Disorder](#)