


CD55 polyclonal antibody

Catalog # PAB30951 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lung with CD55 polyclonal antibody (Cat # PAB30951) shows strong membranous positivity in alveolar cells.

Specification

Product Description	Rabbit polyclonal antibody raised against partial recombinant human CD55.
Immunogen	Recombinant protein corresponding to human CD55.
Sequence	PDVPNAQPALEGRTSFPEDTVITYKCEESFVKIPGEKDSVICLKGSQWSDIEEFCNRSCEVPTRLN SASLKQPYITQNYFPVGTVEYECRPGYRREPSLSPKLTCLQNLKWSTAVEFCKKKSCPNPGEIR NGQIDVPGGILFGATISF
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:50-1:200) The optimal working dilution should be determined by the end user.

Storage Buffer	In PBS, pH 7.2 (40% glycerol, 0.02% sodium azide).
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human lung with CD55 polyclonal antibody (Cat # PAB30951) shows strong membranous positivity in alveolar cells.

Gene Info — CD55

Entrez GeneID	1604
Protein Accession#	P08174
Gene Name	CD55
Gene Alias	CR, CROM, DAF, TC
Gene Description	CD55 molecule, decay accelerating factor for complement (Cromer blood group)
Omim ID	125240
Gene Ontology	Hyperlink
Gene Summary	This gene encodes a protein involved in the regulation of the complement cascade. The encoded glycoprotein is also known as the decay-accelerating factor (DAF); binding of DAF to complement proteins accelerates their decay, disrupting the cascade and preventing damage to host cells. Antigens present on the DAF glycoprotein constitute the Cromer blood group system (CROM). Two alternatively spliced transcripts encoding different proteins have been identified. The predominant transcript encodes a membrane-bound protein expressed on cells exposed to plasma component proteins but an alternatively spliced transcript produces a soluble protein present at much lower levels. Additional, alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq]
Other Designations	CD55 antigen decay accelerating factor for complement

Pathway

- [Complement and coagulation cascades](#)
- [Hematopoietic cell lineage](#)

Disease

- [Asthma](#)
- [Birth Weight](#)
- [Chorioamnionitis](#)
- [Down Syndrome](#)
- [Fetal Membranes](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Leukemia](#)
- [Lymphoma](#)
- [Macular Degeneration](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Obstetric Labor](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Rhinitis](#)