


# Histone H4 (K20me3) polyclonal antibody

Catalog # PAB31330


Size 50 ug

## Applications




### ChIP

ChIP assays were performed using human K562 cells. A titration of the antibody consisting of 0.2, 0.5, 1 and 2 ug per ChIP experiment was analysed. IgG (1 ug/IP) was used as negative IP control. QPCR was performed with primers for promoters of the active genes EIF4A2 and GAPDH, used as negative controls, and for ZNF510 and the Sat2 satellite repeat region used as positive controls. The figure shows the recovery, expressed as a % of input (the relative amount of immunoprecipitated DNA compared to input DNA after qPCR analysis).



### ChIP-Seq

The figure shows the signal distribution along the long arm of chromosome 19 and a zoomin to an enriched region containing several ZNF repeat genes and the enrichment in the telomeric region of chromosome 12, also containing several ZNF repeat genes, and at ZNF510 on chromosome 9, respectively. The position of the amplicon used for ChIP-qPCR is indicated by an arrow.




### Western Blot

Western Blot analysis of (1) 25 ug whole cell extracts of HeLa cells, (2) 15 ug histone extracts of HeLa cells, (3) 1 ug of recombinant histone H2A, (4) 1 ug of recombinant histone H2B, (5) 1 ug of recombinant histone H3, (6) 1 ug of recombinant histone H4.


## Immunofluorescence

Immunofluorescent staining of Hela cell line with antibody followed by an anti-rabbit antibody conjugated to Alexa488 (left). The middle panel shows staining of the nuclei with DAPI. A merge of the two stainings (right).



## Enzyme-linked Immunoabsorbent Assay


ELISA is a quantitative method used to determine the titer of the antibody using a serial dilution of antibody against Histone H4 (K20me3) in antigen coated wells. The antigen used was a peptide containing the histone modification of interest. By plotting the absorbance against the antibody dilution, the titer of the antibody was estimated to be 1:21700.



## Dot Blot

Cross reactivity test using the Histone H4 (K20me3) antibody.

Dot Blot analysis was performed with peptides containing other histone modifications and the unmodified H4K20. One hundred to 0.2 pmol of the respective peptides were spotted on a membrane. The antibody was used at a dilution of 1:20000. The figure shows a high specificity of the antibody for the modification of interest.



## Specification

Product Description	Rabbit polyclonal antibody raised against synthetic peptide of Histone H4 (K20me3).
Immunogen	A synthetic peptide (conjugated with KLH) corresponding to Histone H4, trimethylated at lysine 20.
Host	Rabbit
Reactivity	Human, Mouse
Form	Liquid
Purification	Affinity purification

<b>Recommend Usage</b>	ELISA (1:500) Western Blot (1:1000) ChIP (0.5-1 ug/IP) Dot Blot (1:20000) Immunofluorescence (1:500) The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	In PBS (0.05% sodium azide, 0.05% proclin 300).
<b>Storage Instruction</b>	Store at -20°C. For long term storage store at -80°C. Aliquot to avoid repeated freezing and thawing.
<b>Note</b>	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

### ● ChIP

ChIP assays were performed using human K562 cells. A titration of the antibody consisting of 0.2, 0.5, 1 and 2 ug per ChIP experiment was analysed. IgG (1 ug/IP) was used as negative IP control. QPCR was performed with primers for promoters of the active genes EIF4A2 and GAPDH, used as negative controls, and for ZNF510 and the Sat2 satellite repeat region used as positive controls. The figure shows the recovery, expressed as a % of input (the relative amount of immunoprecipitated DNA compared to input DNA after qPCR analysis).

### ● ChIP-Seq

The figure shows the signal distribution along the long arm of chromosome 19 and a zoom in to an enriched region containing several ZNF repeat genes and the enrichment in the telomeric region of chromosome 12, also containing several ZNF repeat genes, and at ZNF510 on chromosome 9, respectively. The position of the amplicon used for ChIP-qPCR is indicated by an arrow.

### ● Western Blot

Western Blot analysis of (1) 25 ug whole cell extracts of Hela cells, (2) 15 ug histone extracts of Hela cells, (3) 1 ug of recombinant histone H2A, (4) 1 ug of recombinant histone H2B, (5) 1 ug of recombinant histone H3, (6) 1 ug of recombinant histone H4.

### ● Immunofluorescence

Immunofluorescent staining of Hela cell line with antibody followed by an anti-rabbit antibody conjugated to Alexa488 (left). The middle panel shows staining of the nuclei with DAPI. A merge of the two stainings (right).

### ● Enzyme-linked Immunoabsorbent Assay

ELISA is a quantitative method used to determine the titer of the antibody using a serial dilution of antibody against Histone H4 (K20me3) in antigen coated wells. The antigen used was a peptide containing the histone modification of interest. By plotting the absorbance against the antibody dilution, the titer of the antibody was estimated to be 1:21700.

- Dot Blot

Cross reactivity test using the Histone H4 (K20me3) antibody.

Dot Blot analysis was performed with peptides containing other histone modifications and the unmodified H4K20. One hundred to 0.2 pmol of the respective peptides were spotted on a membrane. The antibody was used at a dilution of 1:20000. The figure shows a high specificity of the antibody for the modification of interest.

## Gene Info — HIST1H4A

**Entrez GeneID** [8359](#)

**Protein Accession#** [P62805](#)

**Gene Name** HIST1H4A

**Gene Alias** H4/a, H4FA

**Gene Description** histone cluster 1, H4a

**Omim ID** [602822](#)

**Gene Ontology** [Hyperlink](#)

**Gene Summary** Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by RefSeq]

**Other Designations** H4 histone family, member A|histone 1, H4a

## Pathway

- [Systemic lupus erythematosus](#)