


# DDR1 polyclonal antibody

Catalog # PAB3365

Size 400 uL

## Applications



### Western Blot (Tissue lysate)

Western blot analysis of DDR1 polyclonal antibody (Cat # PAB3365) in placenta lysate. MCK10 (arrow) was detected using purified polyclonal antibody. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed and paraffin-embedded human cancer tissue reacted with DDR1 polyclonal antibody (Cat # PAB3365), which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma.

## Specification

<b>Product Description</b>	Rabbit polyclonal antibody raised against synthetic peptide of DDR1.
<b>Immunogen</b>	A synthetic peptide (conjugated with KLH) corresponding to 24-54 amino acids at N-terminus of human DDR1.
<b>Host</b>	Rabbit
<b>Reactivity</b>	Human
<b>Form</b>	Liquid
<b>Purification</b>	Protein G purification

<b>Recommend Usage</b>	Flow Cytometry (1:10-50) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (1:10-50) Western Blot (1:1000) The optimal working dilution should be determined by the end user.
<b>Storage Buffer</b>	In PBS (0.09% sodium azide)
<b>Storage Instruction</b>	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
<b>Note</b>	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Applications

- Western Blot (Tissue lysate)

Western blot analysis of DDR1 polyclonal antibody (Cat # PAB3365) in placenta lysate. MCK10 (arrow) was detected using purified polyclonal antibody. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Formalin-fixed and paraffin-embedded human cancer tissue reacted with DDR1 polyclonal antibody (Cat # PAB3365), which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma.

- Flow Cytometry

## Gene Info — DDR1

<b>Entrez GeneID</b>	<a href="#">780</a>
<b>Protein Accession#</b>	<a href="#">NP_054699;Q08345</a>
<b>Gene Name</b>	DDR1
<b>Gene Alias</b>	CAK, CD167, DDR, EDDR1, MCK10, NEP, NTRK4, PTK3, PTK3A, RTK6, TRKE
<b>Gene Description</b>	discoidin domain receptor tyrosine kinase 1
<b>Omim ID</b>	<a href="#">600408</a>
<b>Gene Ontology</b>	<a href="#">Hyperlink</a>

## Gene Summary

Receptor tyrosine kinases (RTKs) play a key role in the communication of cells with their microenvironment. These molecules are involved in the regulation of cell growth, differentiation and metabolism. The protein encoded by this gene is a RTK that is widely expressed in normal and transformed epithelial cells and is activated by various types of collagen. This protein belongs to a subfamily of tyrosine kinase receptors with a homology region to the Dictyostelium discoideum protein discoidin I in their extracellular domain. Its autophosphorylation is achieved by all collagens so far tested (type I to type VI). In situ studies and Northern-blot analysis showed that expression of this encoded protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, this protein is significantly over-expressed in several human tumors from breast, ovarian, esophageal, and pediatric brain. This gene is located on chromosome 6p21.3 in proximity to several HLA class I genes. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq]

## Other Designations

OTTHUMP00000029343|OTTHUMP00000029344|OTTHUMP00000029345|OTTHUMP00000029346|OTTHUMP00000029347|PTK3A protein tyrosine kinase 3A|cell adhesion kinase|discoidin domain receptor DDR1d|discoidin domain receptor family, member 1|discoidin receptor tyrosine kinase

## Publication Reference

- [The discoidin domain receptor tyrosine kinases are activated by collagen.](#)

Vogel W, Gish GD, Alves F, Pawson T.

Molecular Cell 1997 Dec; 1(1):13.

Application: WB-Ce, WB-Tr, Human, HEK 293, T47D cells

- [The genomic structure of discoidin receptor tyrosine kinase.](#)

Playford MP, Butler RJ, Wang XC, Katso RM, Cooke IE, Ganesan TS.

Genome Research 1996 Jul; 6(7):620.

Application: WB-Ce, WB-Tr, Human, Mammalian cells

## Disease

- [Abortion](#)
- [Arthritis](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Genetic Predisposition to Disease](#)
- [Glomerulonephritis](#)

- [Leukemia](#)
- [Lupus Erythematosus](#)
- [Schizophrenia](#)
- [Vitiligo](#)