PARK7 polyclonal antibody

Catalog # PAB4487 Size 400 uL

Applications



Western Blot

Western blot analysis of PARK7 polyclonal antibody (Cat # PAB4487) in mouse brain,kidney,liver tissue lysates and NIH-3T3, HeLa cell line lysates (35 ug/lane). PARK7 (arrow) was detected using the purified PARK7 polyclonal antibody (Cat # PAB4487).

Specification	
Product Description	Rabbit polyclonal antibody raised against synthetic peptide of PARK7.
Immunogen	A synthetic peptide (conjugated with KLH) corresponding to N-terminus of human PARK7.
Host	Rabbit
Reactivity	Human
Form	Liquid
Purification	Ammonium sulfate precipitation
Recommend Usage	ELISA (1:1000) Western Blot (1:100-500) The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS (0.09% sodium azide)
Storage Instruction	Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.



Applications

Western Blot

Western blot analysis of PARK7 polyclonal antibody (Cat # PAB4487) in mouse brain,kidney,liver tissue lysates and NIH-3T3, HeLa cell line lysates (35 ug/lane). PARK7 (arrow) was detected using the purified PARK7 polyclonal antibody (Cat # PAB4487).

Enzyme-linked Immunoabsorbent Assay

Gene Info — PARK7	
Entrez GenelD	<u>11315</u>
Protein Accession#	<u>NP_009193;Q99497</u>
Gene Name	PARK7
Gene Alias	DJ-1, DJ1, FLJ27376, FLJ34360, FLJ92274
Gene Description	Parkinson disease (autosomal recessive, early onset) 7
Omim ID	<u>602533</u> <u>606324</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The product of this gene belongs to the peptidase C56 family of proteins. It acts as a positive reg ulator of androgen receptor-dependent transcription. It may also function as a redox-sensitive cha perone, as a sensor for oxidative stress, and it apparently protects neurons against oxidative stre ss and cell death. Defects in this gene are the cause of autosomal recessive early-onset Parkinso n disease 7. Two transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq
Other Designations	OTTHUMP0000001348 OTTHUMP0000001349 OTTHUMP0000001350 OTTHUMP000000 01351 Parkinson disease protein 7 oncogene DJ1 protein DJ-1

Publication Reference

DJ-1 restores p53 transcription activity inhibited by Topors/p53BP3.

Shinbo Y, Taira T, Niki T, Iguchi-Ariga SM, Ariga H.

International Journal of Oncology 2005 Mar; 26(3):641.

Application: IF, WB-Tr, Human, HEK 293T, HeLa, NCI-H1299 cells

😵 Abnova

Product Information

• DJ-1, a novel regulator of the tumor suppressor PTEN.

Kim RH, Peters M, Jang Y, Shi W, Pintilie M, Fletcher GC, DeLuca C, Liepa J, Zhou L, Snow B, Binari RC, Manoukian AS, Bray MR, Liu FF, Tsao MS, Mak TW.

Cancer Cell 2005 Mar; 7(3):263.

Application: IHC-P, WB, Human, A-549 cells, Human breast cancer

Reduced anti-oxidative stress activities of DJ-1 mutants found in Parkinson's disease patients.

Takahashi-Niki K, Niki T, Taira T, Iguchi-Ariga SM, Ariga H.

Biochemical and Biophysical Research Communications 2004 Jul; 320(2):389.

Disease

- Alzheimer disease
- <u>Chromosome Disorders</u>
- Dementia
- Disease Progression
- Genetic Predisposition to Disease
- Lewy Body Disease
- Parkinson disease
- Parkinsonian Disorders