

HTATIP2 polyclonal antibody

Catalog # PAB6602 Size 100 ug

Applications



Western Blot (Tissue lysate)

HTATIP2 polyclonal antibody (Cat # PAB6602) staining (1 ug/mL) of human lung lysate (RIPA buffer, 35 ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Specification	
Product Description	Goat polyclonal antibody raised against synthetic peptide of HTATIP2.
Immunogen	A synthetic peptide corresponding to human HTATIP2.
Sequence	AETEALSKLREDFR-C
Host	Goat
Theoretical MW (kDa)	30.1, 27.0, 14.9
Reactivity	Human
Form	Liquid
Purification	Antigen affinity purification
Concentration	0.5 mg/mL
Quality Control Testing	Antibody Reactive Against Synthetic Peptide.
Recommend Usage	ELISA (1:32000) Western Blot (1-3 ug/mL) The optimal working dilution should be determined by the end user.



Product Information

Storage Buffer	In Tris saline, pH 7.3 (0.5% BSA, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot (Tissue lysate)

HTATIP2 polyclonal antibody (Cat # PAB6602) staining (1 ug/mL) of human lung lysate (RIPA buffer, 35 ug total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

Enzyme-linked Immunoabsorbent Assay

Gene Info — HTATIP2	
Entrez GeneID	10553
Protein Accession#	NP_006401
Gene Name	HTATIP2
Gene Alias	CC3, FLJ26963, SDR44U1, TIP30
Gene Description	HIV-1 Tat interactive protein 2, 30kDa
Omim ID	605628
Gene Ontology	<u>Hyperlink</u>
Gene Summary	0
Other Designations	Tat-interacting protein (30kD) short chain dehydrogenase/reductase family 44U, member 1

Publication Reference

A link between metastasis and resistance to apoptosis of variant small cell lung carcinoma.

Shtivelman E.

Oncogene 1997 May; 14(18):2167.