

LITAF polyclonal antibody

Catalog # PAB7572

Size 100 ug

Specification

Product Description	Goat polyclonal antibody raised against synthetic peptide of LITAF.
Immunogen	A synthetic peptide corresponding to human LITAF.
Sequence	C-PDGKGMNPPSYTQ
Host	Goat
Theoretical MW (kDa)	17.1, 15.8
Form	Liquid
Purification	Antigen affinity purification
Concentration	0.5 mg/mL
Quality Control Testing	Antibody Reactive Against Synthetic Peptide.
Recommend Usage	ELISA (1:8000) The optimal working dilution should be determined by the end user.
Storage Buffer	In Tris saline, pH 7.3 (0.5% BSA, 0.02% sodium azide)
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Enzyme-linked Immunoabsorbent Assay

Gene Info — LITAF

Entrez GeneID	9516
Protein Accession#	NP_004853.2
Gene Name	LITAF
Gene Alias	FLJ38636, MGC116698, MGC116700, MGC116701, MGC125274, MGC125275, MGC125276, PIG7, SIMPLE, TP53I7
Gene Description	lipopolysaccharide-induced TNF factor
Omim ID	601098 603795
Gene Ontology	Hyperlink
Gene Summary	Lipopolysaccharide is a potent stimulator of monocytes and macrophages, causing secretion of tumor necrosis factor-alpha (TNF-alpha) and other inflammatory mediators. This gene encodes lipopolysaccharide-induced TNF-alpha factor, which is a DNA-binding protein and can mediate the TNF-alpha expression by direct binding to the promoter region of the TNF-alpha gene. The transcription of this gene is induced by tumor suppressor p53 and has been implicated in the p53-induced apoptotic pathway. Mutations in this gene cause Charcot-Marie-Tooth disease type 1C (CMT1C) and may be involved in the carcinogenesis of extramammary Paget's disease (EMPD). Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq]
Other Designations	LPS-induced TNF-alpha factor lipopolysaccharide-induced TNF-alpha factor lipopolysaccharide-induced tumor necrosis factor-alpha factor p53-induced gene 7 protein small integral membrane protein of lysosome/late endosome tumor protein p53 inducible protein

Publication Reference

- [LPS-induced TNF-alpha factor \(LITAF\)-deficient mice express reduced LPS-induced cytokine: Evidence for LITAF-dependent LPS signaling pathways.](#)

Tang X, Metzger D, Leeman S, Amar S.

PNAS 2006 Sep; 103(37):13777.

Application: WB, Human, Mouse, Macrophages, Monocytes

Disease

- [Arrhythmias](#)
- [Charcot-Marie-Tooth Disease](#)

- [Death](#)
- [Genetic Predisposition to Disease](#)
- [Lung Neoplasms](#)
- [Multiple Sclerosis](#)
- [Pulmonary Disease](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)