


RecomAb™

# CD209 recombinant monoclonal antibody, clone C209/2749R

Catalog # RAB00488

Size 100 ug

## Applications



### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human small intestine with CD209 recombinant monoclonal antibody, clone C209/2749R (Cat # RAB00488).

## Specification

|                             |                                                                                                                                                          |
|-----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Product Description</b>  | Rabbit recombinant monoclonal antibody raised against full length recombinant human CD209.                                                               |
| <b>Antibody Species</b>     | Rabbit                                                                                                                                                   |
| <b>Immunogen</b>            | Original antibody is raised against recombinant protein corresponding to full length human CD209.                                                        |
| <b>Theoretical MW (kDa)</b> | 45                                                                                                                                                       |
| <b>Reactivity</b>           | Human                                                                                                                                                    |
| <b>Form</b>                 | Liquid                                                                                                                                                   |
| <b>Purification</b>         | Protein A/G purification                                                                                                                                 |
| <b>Isotype</b>              | IgG                                                                                                                                                      |
| <b>Recommend Usage</b>      | Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) (0.5-1 ug/mL)<br>The optimal working dilution should be determined by the end user. |
| <b>Storage Buffer</b>       | In 10 mM PBS                                                                                                                                             |

**Storage Instruction**

Store at -20 to -80°C.  
Aliquot to avoid repeated freezing and thawing.

## Applications

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human small intestine with CD209 recombinant monoclonal antibody, clone C209/2749R (Cat # RAB00488).

## Gene Info — CD209

**Entrez GeneID** [30835](#)

**Protein Accession#** [Q9NNX6](#)

**Gene Name** CD209

**Gene Alias** CDSIGN, CLEC4L, DC-SIGN, DC-SIGN1, MGC129965

**Gene Description** CD209 molecule

**Omim ID** [604672](#) [607948](#) [609423](#)

**Gene Ontology** [Hyperlink](#)

**Gene Summary**

This gene encodes a transmembrane receptor and is often referred to as DC-SIGN because of its expression on the surface of dendritic cells and macrophages. The encoded protein is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses with a large impact on public health. The protein is organized into three distinct domains: an N-terminal transmembrane domain, a tandem-repeat neck domain and C-type lectin carbohydrate recognition domain. The extracellular region consisting of the C-type lectin and neck domains has a dual function as a pathogen recognition receptor and a cell adhesion receptor by binding carbohydrate ligands on the surface of microbes and endogenous cells. The neck region is important for homo-oligomerization which allows the receptor to bind multivalent ligands with high avidity. Variations in the number of 23 amino acid repeats in the neck domain of this protein are rare but have a significant impact on ligand binding ability. This gene is closely related in terms of both sequence and function to a neighboring gene (GeneID 10332; often referred to as L-SIGN). DC-SIGN and L-SIGN differ in their ligand-binding properties and distribution. Alternative splicing results in multiple variants

**Other Designations**

C-type lectin domain family 4, member L|CD209 antigen|Dendritic cell-specific intracellular adhesion molecules (ICAM)-3 grabbing non-integrin

## Disease

- [Acquired Immunodeficiency Syndrome](#)
- [Arthritis](#)
- [Asthma](#)
- [Birth Weight](#)
- [Bronchiolitis](#)
- [Celiac Disease](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Cytomegalovirus Infections](#)
- [Dengue](#)
- [Disease Progression](#)
- [Disease Susceptibility](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Hemophilia A](#)
- [Hepatitis C](#)
- [HIV Infections](#)
- [HTLV-I Infections](#)
- [Hypersensitivity](#)
- [Infant](#)
- [Leukemia](#)
- [Liver Cirrhosis](#)
- [Liver Diseases](#)
- [Lymphadenitis](#)

- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Mycobacterium Infections](#)
- [Nasopharyngeal Neoplasms](#)
- [Periodontitis](#)
- [Respiratory Syncytial Virus Infections](#)
- [Severe Acute Respiratory Syndrome](#)
- [Sexually Transmitted Diseases](#)
- [Tuberculosis](#)