


RecomAb™

ITGA2B recombinant monoclonal antibody, clone 7E3 (Abciximab)

Catalog # RAB01147 Size 200 ug

Applications



Flow Cytometry

Flow-cytometry were stained with an isotype control at a concentration of 1 ug/mL for 30 mins at RT. After washing, bound antibody was detected using a AF488 conjugated donkey anti-rabbit antibody and cells analysed on a FACSCanto flow-cytometer.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human CD41.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against CD41 is the integrin alpha chain 2b forming a fibrinogen receptor in heterodimeric complex with beta 3.
Reactivity	Human
Specificity	Recognises human CD41.
Form	Liquid
Purification	Protein A affinity purification
Isotype	IgG, Kappa

Recommend Usage

Flow cytometry
The optimal working dilution should be determined by the end user.

Storage Buffer

In PBS with 0.02% Proclin 300

Storage Instruction

Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Applications

- Flow Cytometry

Flow-cytometry were stained with an isotype control at a concentration of 1 ug/mL for 30 mins at RT. After washing, bound antibody was detected using a AF488 conjugated donkey anti-rabbit antibody and cells analysed on a FACSCanto flow-cytometer.

Gene Info — ITGA2B

Entrez GeneID

[3674](#)

Protein Accession#

[P08514](#)

Gene Name

ITGA2B

Gene Alias

CD41, CD41B, GP2B, GPIIb, GTA, HPA3

Gene Description

integrin, alpha 2b (platelet glycoprotein IIb of IIb/IIIa complex, antigen CD41)

Omim ID

[273800](#) [607759](#)

Gene Ontology

[Hyperlink](#)

Gene Summary

ITGA2B encodes integrin alpha chain 2b. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. Alpha chain 2b undergoes post-translational cleavage to yield disulfide-linked light and heavy chains that join with beta 3 to form a fibronectin receptor or expressed in platelets that plays a crucial role in coagulation. Mutations that interfere with this role result in thrombasthenia. In addition to adhesion, integrins are known to participate in cell-surface mediated signalling. [provided by RefSeq]

Other Designations

integrin alpha 2b|platelet fibrinogen receptor, alpha subunit|platelet-specific antigen BAK

Pathway

- [Arrhythmogenic right ventricular cardiomyopathy \(ARVC\)](#)
- [ECM-receptor interaction](#)

- [Focal adhesion](#)
- [Hematopoietic cell lineage](#)
- [Hypertrophic cardiomyopathy \(HCM\)](#)
- [Pathways in cancer](#)
- [Regulation of actin cytoskeleton](#)
- [Small cell lung cancer](#)

Disease

- [Abortion](#)
- [Acute Chest Syndrome](#)
- [Acute Disease](#)
- [Alzheimer disease](#)
- [Anemia](#)
- [Angina](#)
- [Antiphospholipid Syndrome](#)
- [Arterial Occlusive Diseases](#)
- [Arteriosclerosis](#)
- [Atherosclerosis](#)
- [Brain Ischemia](#)
- [Cardiovascular Diseases](#)
- [Carotid Stenosis](#)
- [Cerebrovascular Accident](#)
- [Cerebrovascular Disorders](#)
- [Coronary Artery Disease](#)
- [Coronary Disease](#)
- [Coronary Restenosis](#)

- [Coronary Stenosis](#)
- [Coronary Thrombosis](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Diabetic Nephropathies](#)
- [Diabetic Retinopathy](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Graft Occlusion](#)
- [Hemorrhage](#)
- [Hypercholesterolemia](#)
- [Hypertension](#)
- [Infant](#)
- [Inflammation](#)
- [Insulin Resistance](#)
- [Intracranial Hemorrhages](#)
- [Ischemic Attack](#)
- [Kidney Failure](#)
- [Lupus Erythematosus](#)
- [Metabolic Syndrome X](#)
- [Myocardial Infarction](#)
- [Myocardial Ischemia](#)
- [Neoplasms](#)
- [Obesity](#)
- [Osteoporosis](#)
- [Pain](#)

- [Postoperative Complications](#)
- [Pre-Eclampsia](#)
- [Prosthesis Failure](#)
- [Recurrence](#)
- [Retinal Vein Occlusion](#)
- [Stroke](#)
- [Thrombasthenia](#)
- [Thromboangiitis Obliterans](#)
- [Thrombocytopenia](#)
- [Thrombophilia](#)
- [Thrombosis](#)
- [Vascular Diseases](#)