


 $\textbf{RecomAb}^{\text{\tiny{TM}}}$

ITGA4 recombinant monoclonal antibody, clone Natalizumab

Catalog # RAB01173 Size 200 ug

Applications



Flow Cytometry

Flow-cytometry were stained with an isotype control at a concentration of 1 ug/mL for 30 mins at RT. After washing, bound antibody was detected using a AF488 conjugated donkey anti-rabbit antibody and cells analysed on a FACSCanto flow-cytometer.

Specification	
Product Description	Mouse recombinant monoclonal antibody raised against human Integrin alpha 4.
Antibody Species	Mouse
Immunogen	Original antibody is raised against the RAMOS cell line, a non-EBV releasing cell line, injected into mice.
Reactivity	Human
Specificity	This antibody binds to the alpha 4 binds subunit of $\alpha4\beta1$ (also known as very late antigen 4 [VLA-4] o r CD49d-CD29) and $\alpha4\beta7$ integrin.
Form	Liquid
Purification	Protein A affinity purification
Isotype	lgG1, Kappa



Product Information

Recommend Usage	Flow cytometry The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS with 0.02% Proclin 300
Storage Instruction	Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Applications

Flow Cytometry

Flow-cytometry were stained with an isotype control at a concentration of 1 ug/mL for 30 mins at RT. After washing, bound antibody was detected using a AF488 conjugated donkey anti-rabbit antibody and cells analysed on a FACSCanto flow-cytometer.

Gene Info — ITGA4	
Entrez GenelD	<u>3676</u>
Protein Accession#	<u>P13612</u>
Gene Name	ITGA4
Gene Alias	CD49D, IA4, MGC90518
Gene Description	integrin, alpha 4 (antigen CD49D, alpha 4 subunit of VLA-4 receptor)
Omim ID	<u>192975</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	The product of this gene belongs to the integrin alpha chain family of proteins. Integrins are hetero dimeric integral membrane proteins composed of an alpha chain and a beta chain. This gene enc odes an alpha 4 chain. Unlike other integrin alpha chains, alpha 4 neither contains an I-domain, no rundergoes disulfide-linked cleavage. Alpha 4 chain associates with either beta 1 chain or beta 7 chain. [provided by RefSeq
Other Designations	269C wild type antigen CD49D, alpha-4 subunit of VLA-4 receptor integrin alpha 4 integrin alpha-4 subunit very late activation protein 4 receptor, alpha 4 subunit

Pathway

- Arrhythmogenic right ventricular cardiomyopathy (ARVC)
- Cell adhesion molecules (CAMs)



- ECM-receptor interaction
- Focal adhesion
- Hematopoietic cell lineage
- Hypertrophic cardiomyopathy (HCM)
- Leukocyte transendothelial migration
- Regulation of actin cytoskeleton

Disease

- Autistic Disorder
- Cardiovascular Diseases
- Diabetes Mellitus
- Edema
- Genetic Predisposition to Disease
- Graft vs Host Disease
- Multiple Sclerosis
- Neurogenic Inflammation
- Tobacco Use Disorder