


RecomAb™

STAT4 recombinant monoclonal antibody, clone R02-1C1


Catalog # RAB01353 Size 100 uL

Applications



Western Blot

Western blot analysis of STAT4 in HeLa, A549 lysates using STAT4 antibody.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemistry analysis of paraffin-embedded Human lung cancer using STAT4 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human STAT4.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against recombinant protein corresponding to human STAT4.
Theoretical MW (kDa)	Calculated MW: 86 kD
Reactivity	Human
Form	Liquid

Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) Immunoprecipitation Western Blot The optimal working dilution should be determined by the end user.
Storage Buffer	In 50mM Tris-Glycine, pH 7.4, (0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)
Storage Instruction	Store at 4°C. For longer storage, aliquot and store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot

Western blot analysis of STAT4 in Hela, A549 lysates using STAT4 antibody.

- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemistry analysis of paraffin-embedded Human lung cancer using STAT4 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

- Immunoprecipitation

Gene Info — STAT4

Entrez GeneID	6775
Protein Accession#	Q14765
Gene Name	STAT4
Gene Alias	SLEB11
Gene Description	signal transducer and activator of transcription 4
Omim ID	152700 180300 600558
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is essential for mediating responses to IL12 in lymphocytes, and regulating the differentiation of T helper cells. [provided by RefSeq]

Other Designations

OTTHUMP00000163553

Pathway

- [Jak-STAT signaling pathway](#)

Disease

- [Antiphospholipid Syndrome](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Autoimmune Diseases](#)
- [Behcet Syndrome](#)
- [Birth Weight](#)
- [Brain Ischemia](#)
- [Bronchiolitis](#)
- [Cardiovascular Diseases](#)
- [Colitis](#)
- [Crohn Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Endometriosis](#)
- [Genetic Predisposition to Disease](#)
- [Giant Cell Arteritis](#)

- [Glioblastoma](#)
- [Glioma](#)
- [Glomerulonephritis](#)
- [Infant](#)
- [Inflammation](#)
- [Inflammatory Bowel Diseases](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Lupus Erythematosus](#)
- [Lupus Nephritis](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Multiple Sclerosis](#)
- [Osteoarthritis](#)
- [Proteinuria](#)
- [Psoriasis](#)
- [Pulmonary Fibrosis](#)
- [Respiratory Syncytial Virus Infections](#)
- [Scleroderma](#)
- [Tobacco Use Disorder](#)
- [Uveomeningoencephalitic Syndrome](#)
- [Wegener Granulomatosis](#)