


RecomAb™

CDK4 recombinant monoclonal antibody, clone R04-8I5

Catalog # RAB01515 Size 100 uL

Applications



Western Blot

Western blot analysis of CDK4 in K562, HeLa lysates using human CDK4 recombinant monoclonal antibody, clone R04-8I5 (Cat # RAB01515).

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against synthetic peptide of human Cdk4.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human Cdk4
Theoretical MW (kDa)	Calculated MW: 34 kD
Reactivity	Human
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunofluorescence (1:50-1:200) Immunoprecipitation (1:20) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction

Store at 4°C for short term. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot

Western blot analysis of CDK4 in K562, Hela lysates using human CDK4 recombinant monoclonal antibody, clone R04-8I5 (Cat # RAB01515).

- Immunocytochemistry

- Immunoprecipitation

Gene Info — CDK4

Entrez GeneID [1019](#)

Protein Accession# [P11802](#)

Gene Name CDK4

Gene Alias CMM3, MGC14458, PSK-J3

Gene Description cyclin-dependent kinase 4

Omim ID [123829](#) [609048](#)

Gene Ontology [Hyperlink](#)

Gene Summary

The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is highly similar to the gene products of *S. cerevisiae* cdc28 and *S. pombe* cdc2. It is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression. The activity of this kinase is restricted to the G1-S phase, which is controlled by the regulatory subunits D-type cyclins and CDK inhibitor p16(INK4a). This kinase was shown to be responsible for the phosphorylation of retinoblastoma gene product (Rb). Mutations in this gene as well as in its related proteins including D-type cyclins, p16(INK4a) and Rb were all found to be associated with tumorigenesis of a variety of cancers. Multiple polyadenylation sites of this gene have been reported. [provided by RefSeq]

Other Designations cell division kinase 4|melanoma cutaneous malignant, 3

Pathway

- [Bladder cancer](#)
- [Cell cycle](#)
- [Chronic myeloid leukemia](#)
- [Glioma](#)
- [Melanoma](#)
- [Non-small cell lung cancer](#)
- [p53 signaling pathway](#)
- [Pancreatic cancer](#)
- [Pathways in cancer](#)
- [Small cell lung cancer](#)
- [T cell receptor signaling pathway](#)
- [Tight junction](#)

Disease

- [Adenocarcinoma](#)
- [Birth Weight](#)
- [Brain Neoplasms](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Cardiovascular Diseases](#)
- [Diabetes Mellitus](#)
- [Edema](#)
- [Esophageal Neoplasms](#)
- [Genetic Predisposition to Disease](#)

- [Glioblastoma](#)
- [Head and Neck Neoplasms](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Lung Neoplasms](#)
- [Lymphoma](#)
- [Malignant melanoma](#)
- [Melanoma](#)
- [Mouth Neoplasms](#)
- [Neoplasm Invasiveness](#)
- [Neoplasm Recurrence](#)
- [Neoplasms](#)
- [Obesity](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Pancreatic cancer](#)
- [Pancreatic Neoplasms](#)
- [Precancerous Conditions](#)
- [Pulmonary Disease](#)
- [Retinoblastoma](#)
- [Skin Neoplasms](#)
- [Urinary Bladder Neoplasms](#)
- [Uveal Neoplasms](#)
- [Werner syndrome](#)