


RecomAb™

PRKCA (phospho Ser660) recombinant monoclonal antibody, clone R08-0H5

Catalog # RAB01779 Size 100 uL

Applications



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Paraffin-embedded sections) of human brain with PRKCA (phospho Ser660) recombinant monoclonal antibody, clone R08-0H5 (Cat # RAB01779). High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human PRKCA.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic phosphopeptide corresponding to residues surrounding Ser660 of human PRKCA.
Theoretical MW (kDa)	Calculated MW: 77 kD
Reactivity	Human, Rat
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunohistochemistry (1/50-1/100) Western Blot (1/500-1/1000) The optimal working dilution should be determined by the end user.

Storage Buffer	In 50 mM Tris-Glycine pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)
Storage Instruction	Store at -20 °C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot
- Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Paraffin-embedded sections) of human brain with PRKCA (phospho Ser660) recombinant monoclonal antibody, clone R08-0H5 (Cat # RAB01779). High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Gene Info — PRKCA

Entrez GeneID	5578
Protein Accession#	P17252;P05771;Q05655;Q02156;P24723;Q04759
Gene Name	PRKCA
Gene Alias	AAG6, MGC129900, MGC129901, PKC-alpha, PKCA, PRKACA
Gene Description	protein kinase C, alpha
Omim ID	176960
Gene Ontology	Hyperlink
Gene Summary	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca(2+) handling in myocytes. [provided by RefSeq]
Other Designations	aging-associated gene 6 protein kinase C alpha type

Gene Info — PRKCB

Entrez GeneID [5579](#)**Protein Accession#** [P17252;P05771;Q05655;Q02156;P24723;Q04759](#)**Gene Name** PRKCB**Gene Alias** MGC41878, PKC-beta, PKCB, PRKCB1, PRKCB2**Gene Description** protein kinase C, beta**Omim ID** [176970](#)**Gene Ontology** [Hyperlink](#)

Gene Summary

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This protein kinase has been reported to be involved in many different cellular functions, such as B cell activation, apoptosis induction, endothelial cell proliferation, and intestinal sugar absorption. Studies in mice also suggest that this kinase may also regulate neuronal functions and correlate fear-induced conflict behavior after stress. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq]

Other Designations protein kinase C, beta 1 polypeptide

Gene Info — PRKCD

Entrez GeneID [5580](#)**Protein Accession#** [P17252;P05771;Q05655;Q02156;P24723;Q04759](#)**Gene Name** PRKCD**Gene Alias** MAY1, MGC49908, PKCD, nPKC-delta**Gene Description** protein kinase C, delta**Omim ID** [176977](#)**Gene Ontology** [Hyperlink](#)

Gene Summary

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play distinct roles in cells. The protein encoded by this gene is one of the PKC family members. Studies both in human and mice demonstrate that this kinase is involved in B cell signaling and in the regulation of growth, apoptosis, and differentiation of a variety of cell types. Alternatively spliced transcript variants encoding the same protein have been observed. [provided by RefSeq]

Other Designations

protein kinase C delta VIII

Gene Info — PRKCE**Entrez GeneID**[5581](#)**Protein Accession#**[P17252;P05771;Q05655;Q02156;P24723;Q04759](#)**Gene Name**

PRKCE

Gene Alias

MGC125656, MGC125657, PKCE, nPKC-epsilon

Gene Description

protein kinase C, epsilon

Omim ID[176975](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been shown to be involved in many different cellular functions, such as neuron channel activation, apoptosis, cardioprotection from ischemia, heat shock response, as well as insulin exocytosis. Knockout studies in mice suggest that this kinase is important for lipopolysaccharide (LPS)-mediated signaling in activated macrophages and may also play a role in controlling anxiety-like behavior. [provided by RefSeq]

Other Designations

-

Gene Info — PRKCH**Entrez GeneID**[5583](#)**Protein Accession#**[P17252;P05771;Q05655;Q02156;P24723;Q04759](#)

Gene Name	PRKCH
Gene Alias	MGC26269, MGC5363, PKC-L, PKCL, PRKCL, nPKC-eta
Gene Description	protein kinase C, eta
Omim ID	601367 605437
Gene Ontology	Hyperlink
Gene Summary	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. It is a calcium-independent and phospholipids-dependent protein kinase. It is predominantly expressed in epithelial tissues and has been shown to reside specifically in the cell nucleus. This protein kinase can regulate keratinocyte differentiation by activating the MAP kinase MAPK13 (p38delta)-activated protein kinase cascade that targets CCAAT/enhancer-binding protein alpha (CEBPA). It is also found to mediate the transcription activation of the transglutaminase 1 (TGM1) gene. [provided by RefSeq]
Other Designations	protein kinase C eta type

Gene Info — PRKCQ

Entrez GeneID	5588
Protein Accession#	P17252;P05771;Q05655;Q02156;P24723;Q04759
Gene Name	PRKCQ
Gene Alias	MGC126514, MGC141919, PRKCT, nPKC-theta
Gene Description	protein kinase C, theta
Omim ID	600448
Gene Ontology	Hyperlink

Gene Summary

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role. The protein encoded by this gene is one of the PKC family members. It is a calcium-independent and phospholipid-dependent protein kinase. This kinase is important for T-cell activation. It is required for the activation of the transcription factors NF-kappaB and AP-1, and may link the T cell receptor (TCR) signaling complex to the activation of the transcription factors. [provided by RefSeq]

Other Designations

OTTHUMP00000019053|OTTHUMP00000043364|OTTHUMP00000043365

Pathway

- [Adipocytokine signaling pathway](#)
- [B cell receptor signaling pathway](#)
- [Calcium signaling pathway](#)
- [Calcium signaling pathway](#)
- [Chemokine signaling pathway](#)
- [Chemokine signaling pathway](#)
- [ErbB signaling pathway](#)
- [ErbB signaling pathway](#)
- [Fc epsilon RI signaling pathway](#)
- [Fc epsilon RI signaling pathway](#)
- [Fc epsilon RI signaling pathway](#)
- [Fc epsilon RI signaling pathway](#)
- [Fc gamma R-mediated phagocytosis](#)
- [Fc gamma R-mediated phagocytosis](#)
- [Fc gamma R-mediated phagocytosis](#)
- [Fc gamma R-mediated phagocytosis](#)
- [Focal adhesion](#)

- [Focal adhesion](#)
- [Gap junction](#)
- [Gap junction](#)
- [Glioma](#)
- [Glioma](#)
- [GnRH signaling pathway](#)
- [GnRH signaling pathway](#)
- [GnRH signaling pathway](#)
- [Leukocyte transendothelial migration](#)
- [Leukocyte transendothelial migration](#)
- [Long-term depression](#)
- [Long-term depression](#)
- [Long-term potentiation](#)
- [Long-term potentiation](#)
- [MAPK signaling pathway](#)
- [MAPK signaling pathway](#)
- [Melanogenesis](#)
- [Melanogenesis](#)
- [Natural killer cell mediated cytotoxicity](#)
- [Natural killer cell mediated cytotoxicity](#)
- [Neurotrophin signaling pathway](#)
- [Non-small cell lung cancer](#)
- [Non-small cell lung cancer](#)
- [Pathogenic Escherichia coli infection - EHEC](#)
- [Pathways in cancer](#)

- [Pathways in cancer](#)
- [Phosphatidylinositol signaling system](#)
- [Phosphatidylinositol signaling system](#)
- [T cell receptor signaling pathway](#)
- [Tight junction](#)
- [Tight junction](#)
- [Tight junction](#)
- [Tight junction](#)
- [Tight junction](#)
- [Tight junction](#)
- [Type II diabetes mellitus](#)
- [Type II diabetes mellitus](#)
- [Vascular smooth muscle contraction](#)
- [Vascular smooth muscle contraction](#)
- [Vascular smooth muscle contraction](#)
- [Vascular smooth muscle contraction](#)
- [Vascular smooth muscle contraction](#)
- [Vascular smooth muscle contraction](#)
- [VEGF signaling pathway](#)
- [VEGF signaling pathway](#)
- [Vibrio cholerae infection](#)
- [Vibrio cholerae infection](#)
- [Wnt signaling pathway](#)
- [Wnt signaling pathway](#)

Disease

- [Adenocarcinoma](#)

- [Albuminuria](#)
- [Alzheimer Disease](#)
- [Arthritis](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Autistic Disorder](#)
- [Brain Infarction](#)
- [Brain Ischemia](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cardiovascular Diseases](#)
- [Cardiovascular Diseases](#)
- [Celiac Disease](#)
- [Cerebral Hemorrhage](#)
- [Cerebral Hemorrhage](#)
- [Depressive Disorder](#)
- [Diabetes Mellitus](#)
- [Diabetes Mellitus](#)
- [Diabetes Mellitus](#)
- [Diabetic Angiopathies](#)
- [Diabetic Nephropathies](#)
- [Diabetic Retinopathy](#)
- [Disease Models](#)

- [Disease Progression](#)
- [Disease Progression](#)
- [Edema](#)
- [Edema](#)
- [Edema](#)
- [Epilepsies](#)
- [Esophageal Neoplasms](#)
- [Gastritis](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Genetic Predisposition to Disease](#)
- [Helicobacter Infections](#)
- [Hypertension](#)
- [Hypertension](#)
- [Inflammation](#)
- [Inflammation](#)
- [Intracranial Hemorrhages](#)
- [Kidney Failure](#)
- [Kidney Failure](#)
- [Liver Cirrhosis](#)
- [Liver Cirrhosis](#)
- [Mental Disorders](#)

- [Multiple Sclerosis](#)
- [Narcolepsy](#)
- [Narcolepsy](#)
- [Narcolepsy](#)
- [Obesity](#)
- [Pancreatic Neoplasms](#)
- [Premature Birth](#)
- [Prostatic Neoplasms](#)
- [Proteinuria](#)
- [Schizophrenia](#)
- [Stomach Neoplasms](#)
- [Stroke](#)
- [Stroke](#)
- [Subarachnoid Hemorrhage](#)
- [Syndrome](#)
- [Tobacco Use Disorder](#)
- [Tobacco Use Disorder](#)
- [Tobacco Use Disorder](#)
- [Tobacco Use Disorder](#)
- [Tobacco Use Disorder](#)
- [Tobacco Use Disorder](#)
- [Vaginitis](#)
- [Wegener Granulomatosis](#)