


RecomAb™

MMP12 recombinant monoclonal antibody, clone R03-2B9

Catalog # RAB02100 Size 100 uL

Applications



Western Blot

Western Blot analysis of Hela lysates with MMP12 recombinant monoclonal antibody, clone R03-2B9 (Cat # RAB02100).

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human MMP12.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human MMP12.
Theoretical MW (kDa)	Calculated MW: 54 kD
Reactivity	Hamster, Human, Rat
Form	Liquid
Purification	Affinity purification
Isotype	IgG
Recommend Usage	Immunoprecipitation(1:20) Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction

Store at -20 °C.
Aliquot to avoid repeated freezing and thawing.

Note

This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot

Western Blot analysis of Hela lysates with MMP12 recombinant monoclonal antibody, clone R03-2B9 (Cat # RAB02100).

- Immunoprecipitation

Gene Info — MMP12

Entrez GeneID[4321](#)**Protein Accession#**[P39900](#)**Gene Name**

MMP12

Gene Alias

HME, MGC138506, MME

Gene Description

matrix metalloproteinase 12 (macrophage elastase)

Omim ID[601046](#)**Gene Ontology**[Hyperlink](#)**Gene Summary**

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. It is thought that the protein encoded by this gene is cleaved at both ends to yield the active enzyme, but this processing has not been fully described. The enzyme degrades soluble and insoluble elastin. It may play a role in aneurysm formation and studies in mice suggest a role in the development of emphysema. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. [provided by RefSeq]

Other Designations

macrophage elastase|macrophage metalloelastase|matrix metalloproteinase 12|matrix metalloproteinase 12 (macrophage elastase)

Disease

- [Acute Disease](#)
- [Adenocarcinoma](#)
- [alpha 1-Antitrypsin Deficiency](#)
- [Alpha-1-antitrypsin deficiency](#)
- [Amyotrophic lateral sclerosis](#)
- [Anoxia](#)
- [Aortic Aneurysm](#)
- [Arthritis](#)
- [Asthma](#)
- [Atherosclerosis](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Carotid Artery](#)
- [Chorioamnionitis](#)
- [Chronic Disease](#)
- [Colorectal Neoplasms](#)
- [Coronary Aneurysm](#)
- [Coronary Artery Disease](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Edema](#)
- [Endometriosis](#)
- [Esophageal Neoplasms](#)
- [Fetal Membranes](#)

- [Fibrosis](#)
- [Genetic Predisposition to Disease](#)
- [Glomerulonephritis](#)
- [Hepatitis C](#)
- [Kidney Failure](#)
- [Liver Cirrhosis](#)
- [Lung Neoplasms](#)
- [Metabolic Syndrome X](#)
- [Mucocutaneous Lymph Node Syndrome](#)
- [Multiple Sclerosis](#)
- [Nasopharyngeal Neoplasms](#)
- [Neoplasm Invasiveness](#)
- [Neoplasms](#)
- [Obstetric Labor](#)
- [Ovarian cancer](#)
- [Ovarian Neoplasms](#)
- [Periodontitis](#)
- [Pre-Eclampsia](#)
- [Premature Birth](#)
- [Pulmonary Disease](#)
- [Renal Insufficiency](#)
- [Stomach Neoplasms](#)
- [Subarachnoid Hemorrhage](#)
- [Tuberculosis](#)
- [Urinary Bladder Neoplasms](#)

- [Varicose Ulcer](#)
- [Werner syndrome](#)