

RecomAb™

MDH2 recombinant monoclonal antibody, clone R05-4B5

Catalog # RAB02112 Size 100 uL

Applications



Western Blot

Western Blot analysis of Lane 1: C6, Lane 2: 3T3 and Lane 3: Hela lysates with MDH2 recombinant monoclonal antibody, clone R05-4B5 (Cat # RAB02112).



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human breast cancer with MDH2 recombinant monoclonal antibody, clone R05-4B5 (Cat # RAB020112). High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Specification	
Product Description	Rabbit recombinant monoclonal antibody raised against human MDH2.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against a synthetic peptide corresponding to human MDH2.
Theoretical MW (kDa)	Calculated MW: 36 kD
Reactivity	Human, Mouse, Rat
Form	Liquid

😵 Abnova

Product Information

Purification	Affinity purification
lsotype	lgG
Recommend Usage	Immunohistochemistry (1:50-1:100)
	Western Blot (1:500-1:1000)
	The optimal working dilution should be determined by the end user.
Storage Buffer	In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)
Storage Instruction	Store at -20 °C.
	Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Western Blot

Western Blot analysis of Lane 1: C6, Lane 2: 3T3 and Lane 3: Hela lysates with MDH2 recombinant monoclonal antibody, clone R05-4B5 (Cat # RAB02112).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)

Immunohistochemical staining (Formalin-fixed paraffin-embedded sections) of human breast cancer with MDH2 recombinant monoclonal antibody, clone R05-4B5 (Cat # RAB020112). High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Gene Info — MDH2

Entrez GenelD	<u>4191</u>
Protein Accession#	<u>P40926</u>
Gene Name	MDH2
Gene Alias	M-MDH, MDH, MGC:3559, MOR1
Gene Description	malate dehydrogenase 2, NAD (mitochondrial)
Omim ID	<u>154100</u>
Gene Ontology	Hyperlink



Product Information

Gene Summary

Malate dehydrogenase catalyzes the reversible oxidation of malate to oxaloacetate, utilizing the N AD/NADH cofactor system in the citric acid cycle. The protein encoded by this gene is localized t o the mitochondria and may play pivotal roles in the malate-aspartate shuttle that operates in the metabolic coordination between cytosol and mitochondria. [provided by RefSeq

Other Designations

mitochondrial malate dehydrogenase

Pathway

- Biosynthesis of alkaloids derived from histidine and purine
- Biosynthesis of alkaloids derived from ornithine
- Biosynthesis of alkaloids derived from shikimate pathway
- Biosynthesis of alkaloids derived from terpenoid and polyketide
- Biosynthesis of phenylpropanoids
- Biosynthesis of plant hormones
- Biosynthesis of terpenoids and steroids
- <u>Carbon fixation in photosynthetic organisms</u>
- <u>Citrate cycle (TCA cycle)</u>
- <u>Glyoxylate and dicarboxylate metabolism</u>
- Metabolic pathways
- Pyruvate metabolism
- <u>Reductive carboxylate cycle (CO2 fixation)</u>

Disease

<u>Tobacco Use Disorder</u>