


RecomAb™

HMGCR recombinant monoclonal antibody

Catalog # RAB02643 Size 100 uL

Applications



Western Blot (Cell lysate)

Western blot analysis with HMGCR recombinant monoclonal antibody (Cat # RAB02643) at 1:1000 dilution.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human HMGCR.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against KLH conjugated synthetic peptide derived from human HMGCR.
Theoretical MW (kDa)	97
Reactivity	Human
Specificity	This antibody detects endogenous levels of HMGCR protein.
Form	Liquid
Purification	Protein A purification
Isotype	IgG
Recommend Usage	Western Blot (1:500-1:2000) The optimal working dilution should be determined by the end user.
Storage Buffer	In 0.01M TBS, pH7.4 (1% BSA, 0.03% Proclin300 and 50% Glycerol)

Storage Instruction

Store at 4°C short term.
Aliquot and store at -20°C long term.
Avoid freeze-thaw cycles.

Applications

- Western Blot (Cell lysate)

Western blot analysis with HMGCR recombinant monoclonal antibody (Cat # RAB02643) at 1:1000 dilution.

Gene Info — HMGCR

Entrez GeneID [3156](#)

Protein Accession# [P04035](#)

Gene Name HMGCR

Gene Alias -

Gene Description 3-hydroxy-3-methylglutaryl-Coenzyme A reductase

Omim ID [142910](#)

Gene Ontology [Hyperlink](#)

Gene Summary

HMG-CoA reductase is the rate-limiting enzyme for cholesterol synthesis and is regulated via a negative feedback mechanism mediated by sterols and non-sterol metabolites derived from mevalonate, the product of the reaction catalyzed by reductase. Normally in mammalian cells this enzyme is suppressed by cholesterol derived from the internalization and degradation of low density lipoprotein (LDL) via the LDL receptor. Competitive inhibitors of the reductase induce the expression of LDL receptors in the liver, which in turn increases the catabolism of plasma LDL and lowers the plasma concentration of cholesterol, an important determinant of atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations 3-hydroxy-3-methylglutaryl CoA reductase (NADPH)|hydroxymethylglutaryl-CoA reductase

Pathway

- [Biosynthesis of alkaloids derived from terpenoid and polyketide](#)
- [Biosynthesis of plant hormones](#)
- [Biosynthesis of terpenoids and steroids](#)

- [Metabolic pathways](#)
- [Terpenoid backbone biosynthesis](#)

Disease

- [Acute Disease](#)
- [Alzheimer disease](#)
- [Atherosclerosis](#)
- [Calcinosis](#)
- [Carcinoma](#)
- [Cardiovascular Diseases](#)
- [Cognition Disorders](#)
- [Colorectal Neoplasms](#)
- [Coronary Artery Disease](#)
- [Coronary Disease](#)
- [Dementia](#)
- [Diabetes Complications](#)
- [Diabetes Mellitus](#)
- [Disease Progression](#)
- [Diseases in Twins](#)
- [Drug Toxicity](#)
- [Ductus Arteriosus](#)
- [Dyslipidemias](#)
- [Edema](#)
- [Genetic Predisposition to Disease](#)
- [Hypercholesterolemia](#)
- [Hyperlipidemias](#)

- [Hypertension](#)
- [Infant](#)
- [Liver Neoplasms](#)
- [Lung Neoplasms](#)
- [Metabolic Syndrome X](#)
- [Myocardial Infarction](#)
- [Neoplasms](#)
- [Obstetric Labor](#)
- [Osteoporosis](#)
- [Polycystic Ovary Syndrome](#)
- [Pulmonary Disease](#)
- [Stroke](#)
- [Urinary Bladder Neoplasms](#)
- [Werner syndrome](#)