


RecomAb™

STAT6 recombinant monoclonal antibody, clone Stat6Y641-G12


Catalog # RAB02778 Size 200 uL

Applications



Western Blot

Western blot analysis of Daudi cell extract untreated or treated with IL-4 using 0.01ug/mL Phospho-Stat6 (Tyr641) antibody Stat6Y641-G12.



Flow Cytometry

Flow cytometric analysis of NIH3T3 cells secondary antibody only negative control (blue) or 0.1 ug/mL of isotype control (orange) or treated with imatinib (red) or with pervanadate (green) using Phospho-Stat6 (Tyr641) antibody Stat6Y641-G12 at 0.1 ug/mL.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human STAT6.
Antibody Species	Rabbit
Immunogen	A synthetic phospho-peptide corresponding to residues surrounding Tyr641 of human phospho Stat6
Reactivity	Human
Form	Liquid

Purification	Protein A+G
Isotype	Rabbit IgG1k
Recommend Usage	Flow Cytometry Western Blot The optimal working dilution should be determined by the end user.
Storage Buffer	1X PBS, 0.02% Sodium azide, 50% Glycerol, 0.1% BSA
Storage Instruction	Store at -20°C. Aliquot to avoid repeated freezing and thawing.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Western Blot

Western blot analysis of Daudi cell extract untreated or treated with IL-4 using 0.01ug/mL Phospho-Stat6 (Tyr641) antibody Stat6Y641-G12.

- Flow Cytometry

Flow cytometric analysis of NIH3T3 cells secondary antibody only negative control (blue) or 0.1 ug/mL of isotype control (orange) or treated with imatinib (red) or with pervanadate (green) using Phospho-Stat6 (Tyr641) antibody Stat6Y641-G12 at 0.1 ug/mL.

Gene Info — STAT6

Entrez GeneID	6778
Protein Accession#	P42226
Gene Name	STAT6
Gene Alias	D12S1644, IL-4-STAT, STAT6B, STAT6C
Gene Description	signal transducer and activator of transcription 6, interleukin-4 induced
Omim ID	601512
Gene Ontology	Hyperlink

Gene Summary

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein plays a central role in exerting IL4 mediated biological responses. It is found to induce the expression of BCL2L1/BCL-X(L), which is responsible for the anti-apoptotic activity of IL4. Knockout studies in mice suggested the roles of this gene in differentiation of T helper 2 (Th2) cells, expression of cell surface markers, and class switch of immunoglobulins. [provided by RefSeq]

Other Designations

STAT, interleukin4-induced|signal transducer and activator of transcription 6|transcription factor IL-4 STAT

Pathway

- [Jak-STAT signaling pathway](#)

Disease

- [Anaphylaxis](#)
- [Arthritis](#)
- [Ascariasis](#)
- [Asthma](#)
- [Birth Weight](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Bronchial Hyperreactivity](#)
- [Bronchiolitis](#)
- [Cardiovascular Diseases](#)
- [Colitis](#)
- [Confusion](#)
- [Crohn Disease](#)
- [Dermatitis](#)
- [Diabetes Mellitus](#)

- [Disease Progression](#)
- [Drug Hypersensitivity](#)
- [Edema](#)
- [Endometriosis](#)
- [Food Hypersensitivity](#)
- [Genetic Predisposition to Disease](#)
- [Glioblastoma](#)
- [Glioma](#)
- [Graves Disease](#)
- [Hypersensitivity](#)
- [Infant](#)
- [Infertility](#)
- [Inflammatory Bowel Diseases](#)
- [Kidney Failure](#)
- [Leukemia](#)
- [Liver Diseases](#)
- [Lupus Erythematosus](#)
- [Lymphoma](#)
- [Meningeal Neoplasms](#)
- [Meningioma](#)
- [Nausea](#)
- [Neoplasms](#)
- [Nephrosis](#)
- [Nephrotic Syndrome](#)
- [Nut Hypersensitivity](#)

- [Pain](#)
- [Pulmonary Disease](#)
- [Recurrence](#)
- [Respiratory Syncytial Virus Infections](#)