


RecomAb™

PLCG2 recombinant monoclonal antibody, clone PLCG2Y759-G3 (PE)

Catalog # RAB02887

Size 100 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of Ramos cells unstained treated with imatinib (blue) or stained and treated with imatinib (red) or treated with pervanadate (green) using phospho-PLCγ2 (Tyr759) antibody PLCG2Y759-G3 PE conjugate.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human PLCG2.
Antibody Species	Rabbit
Immunogen	A synthetic phospho-peptide corresponding to residues surrounding Tyr759 of human phospho PLC g2.
Reactivity	Human
Form	Liquid
Purification	Protein A+G
Isotype	Rabbit IgG1k
Conjugation Note	PE

Recommend Usage	Flow Cytometry The optimal working dilution should be determined by the end user.
Storage Buffer	1X PBS, 0.09% Sodium azide, 0.2% BSA
Storage Instruction	Store at 4°C. Do not freeze.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of Ramos cells unstained treated with imatinib (blue) or stained and treated with imatinib (red) or treated with pervanadate (green) using phospho-PLCγ2 (Tyr759) antibody PLCG2Y759-G3 PE conjugate.

Gene Info — PLCG2

Entrez GeneID	5336
Protein Accession#	P16885
Gene Name	PLCG2
Gene Alias	-
Gene Description	phospholipase C, gamma 2 (phosphatidylinositol-specific)
Omim ID	600220
Gene Ontology	Hyperlink
Gene Summary	Enzymes of the phospholipase C family catalyze the hydrolysis of phospholipids to yield diacylglycerols and water-soluble phosphorylated derivatives of the lipid head groups. A number of these enzymes have specificity for phosphoinositides. Of the phosphoinositide-specific phospholipase C enzymes, C-beta is regulated by heterotrimeric G protein-coupled receptors, while the closely related C-gamma-1 (PLCG1; MIM 172420) and C-gamma-2 enzymes are controlled by receptor tyrosine kinases. The C-gamma-1 and C-gamma-2 enzymes are composed of phospholipase domains that flank regions of homology to noncatalytic domains of the SRC oncogene product, SH2 and SH3.[supplied by OMIM]
Other Designations	phospholipase C gamma 2 phospholipase C, gamma 2 phospholipase C, gamma 2 (phosphatidylinositol-specific)

Pathway

- [B cell receptor signaling pathway](#)
- [Calcium signaling pathway](#)
- [Epithelial cell signaling in Helicobacter pylori infection](#)
- [ErbB signaling pathway](#)
- [Fc epsilon RI signaling pathway](#)
- [Fc gamma R-mediated phagocytosis](#)
- [Glioma](#)
- [Inositol phosphate metabolism](#)
- [Leukocyte transendothelial migration](#)
- [Metabolic pathways](#)
- [Natural killer cell mediated cytotoxicity](#)
- [Neurotrophin signaling pathway](#)
- [Non-small cell lung cancer](#)
- [Pathways in cancer](#)
- [Phosphatidylinositol signaling system](#)
- [VEGF signaling pathway](#)
- [Vibrio cholerae infection](#)

Disease

- [Bipolar Disorder](#)
- [Breast cancer](#)
- [Breast Neoplasms](#)
- [Genetic Predisposition to Disease](#)
- [HIV Infections](#)

- [Mental Disorders](#)
- [Ovarian cancer](#)
- [Tobacco Use Disorder](#)