


RecomAb™

H2AX recombinant monoclonal antibody, clone HisH2AXS139-1B3 (APC)

Catalog # RAB02970 Size 100 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of 293T cells, untreated and unstained as negative control (blue) or untreated and stained (red) or treated with UV and PMA and stained (green) using Phospho-Histone H2A.X (Ser139) antibody HisH2AXS139-1B3 APC conjugate.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human H2AX.
Antibody Species	Rabbit
Immunogen	A synthetic phospho-peptide corresponding to residues surrounding Ser139 of human phospho-histo ne H2A.X
Reactivity	Human
Form	Liquid
Conjugation	APC
Purification	Protein A purification, Protein G purification
lsotype	lgG
Recommend Usage	Flow Cytometry The optimal working dilution should be determined by the end user.

Copyright © 2023 Abnova Corporation. All Rights Reserved.

😚 Abnova 🛛

Storage Buffer	1X PBS, 0.09% Sodium azide, 0.2% BSA
Storage Instruction	Store at 4°C. Do not freeze.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which shoul d be handled by trained staff only.

Applications

Flow Cytometry

Flow cytometric analysis of 293T cells, untreated and unstained as negative control (blue) or untreated and stained (red) or treated with UV and PMA and stained (green) using Phospho-Histone H2A.X (Ser139) antibody HisH2AXS139-1B3 APC conjugate.

Gene Info — H2AFX

Entrez GenelD	<u>3014</u>
Protein Accession#	<u>P16104</u>
Gene Name	H2AFX
Gene Alias	H2A.X, H2A/X, H2AX
Gene Description	H2A histone family, member X
Omim ID	<u>601772</u>
Gene Ontology	<u>Hyperlink</u>
Gene Summary	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chro mosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, an d H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and f unctions in the compaction of chromatin into higher order structures. This gene encodes a membe r of the histone H2A family, and generates two transcripts through the use of the conserved stem-l oop termination motif, and the polyA addition motif. [provided by RefSeq
Other Designations	H2AX histone

Pathway

• Systemic lupus erythematosus



Disease

- Azoospermia
- Breast cancer
- Breast Neoplasms
- DNA Damage
- Genetic Predisposition to Disease
- Lymphoma
- Oligospermia
- Ovarian cancer
- Prostate cancer
- Prostatic Neoplasms
- Urinary Bladder Neoplasms