


RecomAb™

BLNK recombinant monoclonal antibody, clone BLNKY84-H4 (FITC)

Catalog # RAB03018

Size 100 Reactions

Applications



Flow Cytometry

Flow cytometric analysis of Ramos cells, unstained and untreated cells as negative control (blue) or untreated (red) or treated with INF α +IL-4 + pervanadate (green) and stained using Phospho-BLNK (Tyr84) antibody BLNKY84-H4 FITCconjugate.

Specification

Product Description	Rabbit recombinant monoclonal antibody raised against human BLNK.
Antibody Species	Rabbit
Immunogen	A synthetic phospho-peptide corresponding to residues surrounding Tyr84 of human phospho BLNK
Reactivity	Human
Form	Liquid
Conjugation	FITC
Purification	Protein A purification, Protein G purification
Isotype	IgG
Recommend Usage	Flow Cytometry The optimal working dilution should be determined by the end user.

Storage Buffer	1X PBS, 0.09% Sodium azide, 0.2% BSA
Storage Instruction	Store at 4°C. Do not freeze.
Note	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Applications

- Flow Cytometry

Flow cytometric analysis of Ramos cells, unstained and untreated cells as negative control (blue) or untreated (red) or treated with INF α +IL-4 + pervanadate (green) and stained using Phospho-BLNK (Tyr84) antibody BLNKY84-H4 FITCconjugate.

Gene Info — BLNK

Entrez GeneID	29760
Protein Accession#	Q8WV28
Gene Name	BLNK
Gene Alias	BASH, BLNK-S, LY57, MGC111051, SLP-65, SLP65
Gene Description	B-cell linker
Omim ID	604515
Gene Ontology	Hyperlink

Gene Summary	This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]
--------------	---

Other Designations	B cell linker protein B-cell adapter containing a SH2 domain protein B-cell adapter containing a Src homology 2 domain protein OTTHUMP00000020167 Src homology 2 domain-containing leukocyte protein of 65 kDa
--------------------	--

Pathway

- [B cell receptor signaling pathway](#)
- [Primary immunodeficiency](#)

Disease

- [Alzheimer Disease](#)
- [Genetic Predisposition to Disease](#)
- [Tobacco Use Disorder](#)