


RecomAb™

CD37 recombinant monoclonal antibody, clone IPO-24 (ZC37-24)


Catalog # RAB03239 Size 200 ug

Applications



Western Blot

Western blot analysis of Lane1: Daudi cell lysate, Lane 2: human spleen and Lane 3: tonsil tissue lysate with CD37 recombinant monoclonal antibody, clone IPO-24 (ZC37-24) (Cat # RAB03239).



Immunofluorescence

Immunofluorescent staining of Daudi cells with CD37 recombinant monoclonal antibody, clone IPO-24 (ZC37-24)(Cat # RAB03239).

Immunofluorescence analysis of paraformaldehyde fixed Daudi cells on Shi-fix™ coverslips stained with the chimeric r version of RAB03239 at 10 ug/ml for 1h followed by Alexa Fluor® 488 secondary antibody (2 ug/ml)- showing membrane staining. The nuclear stain is DAPI (blue). Panels show from left-right- top-bottom RAB03239- DAPI- merged channels and an isotype control. The isotype control was an unknown specificity antibody (3.0) followed by staining with Alexa Fluor® 488 secondary antibody.

Specification	
Product Description	Rabbit recombinant monoclonal antibody raised against human CD37.
Antibody Species	Rabbit
Immunogen	Original antibody is raised against spleen cells of a patient with hairy cell leukemia.
Reactivity	Human



Product Information

Form	Liquid
Isotype	lgG
Recommend Usage	Flow Cytometry
	Immunofluorescence
	Immunohistochemistry
	Western Blot
	The optimal working dilution should be determined by the end user.
Storage Buffer	In PBS with 0.02% Proclin 300
Storage Instruction	Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.
	Aliquot to avoid repeated freezing and thawing.

Applications

Western Blot

Western blot analysis of Lane1: Daudi cell lysate, Lane 2: human spleen and Lane 3: tonsil tissue lysate with CD37 recombinant monoclonal antibody, clone IPO-24 (ZC37-24) (Cat # RAB03239).

- Immunohistochemistry
- Immunofluorescence

Immunofluorescent staining of Daudi cells with CD37 recombinant monoclonal antibody, clone IPO-24 (ZC37-24)(Cat # RAB03239).

Immunofluorescence analysis of paraformaldehyde fixed Daudi cells on Shi-fix™ coverslips stained with the chimeric r version of RAB03239 at 10 ug/ml for 1h followed by Alexa Fluor® 488 secondary antibody (2 ug/ml)- showing membrane staining. The nuclear stain is DAPI (blue). Panels show from left-right- top-bottom RAB03239- DAPI- merged channels and an isotype control. The isotype control was an unknown specificity antibody (3.0) followed by staining with Alexa Fluor® 488 secondary antibody.

Flow Cytometry

Gene Info — CD37		
Entrez GeneID	<u>951</u>	
Gene Name	CD37	
Gene Alias	GP52-40, MGC120234, TSPAN26	
Gene Description	CD37 molecule	
Omim ID	<u>151523</u>	



Product Information

Gene Ontology	<u>Hyperlink</u>
Gene Summary	The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known a s the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins and other transmembrane 4 superfamily proteins. It may play a role in T-cell-B-cell interactions. Alternate splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq
Other Designations	CD37 antigen cell differentiation antigen 37 leukocyte surface antigen CD37 tetraspanin-26

Pathway

• Hematopoietic cell lineage

Disease

- Cardiovascular Diseases
- Diabetes Mellitus
- Edema